

The spirochetes include the causative agents for _____

syphilis

Lyme disease

both syphilis and Lyme disease.

neither syphilis nor Lyme disease.

Rickettsia rickettsii is responsible for

Q-fever

Rocky Mountain spotted fever

Lyme disease

Typhoid fever

Helicobacter pylori is responsible for

gastroenteritis.

cholera.

bacterial dysentery.

peptic ulcer disease.

Whooping cough is caused by

Klebsiella pneumoniae

Vibrio parahaemolyticus

Bordetella pertussis

Yersinia pestis

Which of the following is associated with dental caries?

Streptococcus pneumoniae

Streptococcus mutans

Lactococcus lactis

Staphylococcus aureus

Which of the following is responsible for food spoilage even in canned foods?

Clostridium perfringens

Clostridium botulinum

Clostridium tetani

None of the above

Which of the following is a normal skin resident?

Staphylococcus epidermidis

Staphylococcus aureus

Thermoactinomyces vulgaris

None of the above

Spore-forming bacteria are found mainly
in _____

freshwater environments

marine environments

soil environments

None of the above

E. coli O157:H7 is thought to have acquired enterohemorrhagic genes from:

Clostridium

Bacillus

Campylobacter

Shigella

The main reservoir of *Staphylococcus aureus* is _____

human feces.

human nasal cavity.

human red blood cells.

All of the above.