



Metabolism of lipids II:

Synthesis of fatty acids

Prof. Mamoun Ahram

Resources



- This lecture
- Lippincott's Biochemistry, Ch. 16


Overview of fatty acid synthesis



**Major sites: liver
and adipose tissues**

- The fatty acids are synthesized by:
 1. Production of malonyl CoA
 2. Binding of acetyl CoA and malonyl CoA to the fatty acid synthase
 3. Condensation of acetyl CoA and malonyl CoA
 4. Elongation of the acyl CoA by 2 carbons per round
 - Reduction, dehydration, reduction
 5. Binding of malonyl CoA
 6. *Repeat 3 (acyl CoA), 4, and 5*
 7. Release of the hydrocarbon chain by a thioesterase (TE)

Mitochondria to cytoplasm transport of acetyl-CoA



When ATP increases:

ATP inhibits isocitrate dehydrogenase

Citrate is transported into the cytosol

Citrate is cleaved into oxaloacetate and acetyl CoA by ATP citrate lyase






Figure 21-10
Lehninger Principles of Biochemistry, Fifth Edition
© 2008 W. H. Freeman and Company

Glucose can be converted to fat, but fat cannot be converted to glucose.

Synthesis of malonyl-CoA





- Acetyl CoA carboxylase (ACC) transfers a carbon from CO_2 (as a bicarbonate) via biotin (vitamin B7), which is covalently bound to a lysyl residue of the ACC.
- ATP is needed.
- The reaction is a rate-limiting reaction.
- ACC is an allosteric enzyme.



Regulation of ACC





- ACC is inactivated by:
 - Palmitoyl-CoA
 - Phosphorylation by AMPK, which is activated by glucagon and epinephrine.



Regulation of ACC synthesis




- ACC synthesis is regulated transcription factors:
 - The carbohydrate response element-binding protein (ChREBP)
 - ChREBP is inactivated by phosphorylation by PKA and AMPK preventing its nuclear localization.
 - It is dephosphorylated by excess glucose.
 - The sterol regulatory element-binding protein-1c (SREBP-1c)
 - SREBP-1 is activated by Insulin.
- Fatty acid synthase, glucokinase, ATP citrate lyase and liver pyruvate kinase are similarly regulated.



Metformin



- Metformin lowers plasma TAG through
 - Activation of AMPK, resulting in inhibition of ACC activity (by phosphorylation) and inhibition of ACC and fatty acid synthase expression (by decreasing ChREBP and SREBP-1c).
 - It lowers blood glucose by increasing AMPK-mediated glucose uptake by muscle.








metformin


Fatty acid synthase (FAS)



- A multifunctional, homodimeric enzyme
- Each FAS monomer is multicatalytic with six enzymic domains and a domain for binding a phosphopantetheine-containing acyl carrier protein (ACP) domain.
- Phosphopantetheine, a derivative of pantothenic acid (vitamin B5), carries acyl units on its terminal thiol (–SH) group and presents them to the catalytic domains of FAS.
- It also is a component of CoA.












Condensation, reduction, dehydration, reduction




[4]

CO₂


3-Ketoacyl-ACP synthase
(condensing enzyme)




3-Ketoacyl-ACP reductase




3-Hydroxyacyl-ACP dehydratase



Enoyl-ACP reductase



Compare to the first reaction in the previous slide




The lactating mammary gland terminates lengthening the chain EARLY.


The stoichiometry of palmitate synthesis




- **Stoichiometry of palmitate synthesis:**



- **Malonyl-CoA synthesis:**




- **Overall stoichiometry of palmitate synthesis:**



Sources of molecules




- Acetyl CoA
 - Pyruvate
- NADH (for oxaloacetate to malate)
 - Glycolysis
- NADPH:
 - Pentose phosphate pathway
 - Malate to pyruvate



Further elongation




- Location: smooth endoplasmic reticulum
- Different enzymes are needed.
- Two-carbon donor: Malonyl CoA
- Source of electrons: NADPH
- No ACP or multifunctional enzyme is needed.



- Note: the brain has additional enzymes allowing it to produce the very-long-chain fatty acids ([VLCFA] over 22 carbons)




- Location: mitochondria
- Two-carbon donor: Acetyl CoA
- Source of electrons: NADPH and NADH
- Substrates: fatty acids shorter than 16



Chain desaturation



- Enzymes: fatty acyl CoA desaturases
- Substrates: long-chain fatty acids
- Location: smooth endoplasmic reticulum
- Acceptor of electrons: oxygen (O_2), cytochrome b5, and its flavin adenine dinucleotide (FAD)-linked reductase
- Donor of electrons: NADH
- The first double bond is inserted between carbons 9 and 10, producing oleic acid, 18:1(9), and small amounts of palmitoleic acid, 16:1(9).






Humans have carbon 9, 6, 5, and 4 desaturases but cannot introduce double bonds from carbon 10 to the ω end of the chain. Therefore, the polyunsaturated ω -6 linoleic acid and ω -3 linolenic acid are essential.


Triacylglycerol structure and synthesis



- The fatty acid on carbon 1 is typically saturated, that on carbon 2 is typically unsaturated, and that on carbon 3 can be either.
- Synthesis involves three steps:
 - Glycerol 3-phosphate synthesis
 - Liver (2 mechanisms) vs. adipose tissue (one mechanism only)
 - Activation of fatty acids
 - Synthesis of triacylglycerol



Synthesis of triacylglycerol



Intestinal mucosal cells



- In addition to these two pathways (as in the liver), TAG is synthesized via the MAG pathway in the intestinal mucosal cells.

