

Cardiac output and Venous Return

Faisal I. Mohammed, MD, PhD


Yanal A. Shafagoj MD, PhD

Objectives

- Define cardiac output and venous return
- Describe the methods of measurement of CO
- Outline the factors that regulate cardiac output
- Follow up the cardiac output curves at different physiological states
- Define venous return and describe venous return curve
- Outline the factors that regulate venous return curve at different physiological states
- Inter-relate Cardiac output and venous return curves

Important Concepts About Cardiac Output (CO) Control


- Cardiac Output is the sum of all tissue flows and is affected by their regulation (CO = 5L/min, cardiac index = 3L/min/m²).
- CO is proportional to tissue O₂ use.
- CO is proportional to 1/TPR when AP is constant.
- **$F = \Delta P / R$** (Ohm's law)
- **$CO = (MAP - RAP) / TPR$, ($RAP = 0$) then**
- **$CO = MAP / TPR$; $MAP = CO * TPR$**



Blood flow (ml/g/min)

Tissue	Blood flow (ml/g/min)	A-V difference (Vol %)	Flow ml/min	O ₂ consumption ml/min
Heart	0.8	11	250	27
Brain	0.5	6.2 (25-30% Extraction)	750-900	
Skeletal Muscle	0.03	6	1200	70
Liver	0.6	3.4 Reconditioner organ		
SKIN	0.1			
Kidney	4.2	1.4	1250	18
Carotid bodies	20	0.5	0.6	


Magnitude & Distribution of CO at Rest & During Moderate Exercise




Variations in Tissue Blood Flow

	Per cent	ml/min	ml/min/ 100 gm
Brain	14	700	50
Heart	4	200	70
Bronchi	2	100	25
Kidneys	22	1100	360
Liver	27	1350	95
Portal	(21)	(1050)	
Arterial	(6)	(300)	
Muscle (inactive state)	15	750	4
Bone	5	250	3
Skin (cool weather)	6	300	3
Thyroid gland	1	50	160
Adrenal glands	0.5	25	300
Other tissues	3.5	175	1.3
Total	100.0	5000	---



Control of Cardiac Output




Factors that affect the Cardiac Output




Ventricular Stroke Work Output




CARDIAC OUTPUT CURVES



Effect of Sympathetic and Parasympathetic Stimulation on Cardiac Output



IPP = INTRAPLEURAL PRESSURE




The Cardiac Output Curve

- Plateau of CO curve determined by heart strength (contractility + HR)
- Sympathetics \Rightarrow plateau
- \downarrow Parasympathetics (HR) \Rightarrow (? plateau)
- Plateau
- Heart hypertrophy's \Rightarrow plateau
- Myocardial infarction \Rightarrow (? plateau)
- \downarrow Plateau


The Cardiac Output Curve (cont'd)

- Valvular disease \Rightarrow \downarrow plateau
(stenosis or regurgitation)
- Myocarditis \Rightarrow \downarrow plateau
- Cardiac tamponade \Rightarrow (? plateau)
- \downarrow Plateau
- Metabolic damage \Rightarrow \downarrow plateau


Factors Affecting Cardiac Output




Factors Affecting Stroke Volume




A Summary of the Factors Affecting Cardiac Output



REGULATION OF STROKE VOLUME: PRELOAD




REGULATION OF STROKE VOLUME: CONTRACTILITY



Cardiac Contractility

- Best is to measure the C.O. curve, but this is nearly impossible in humans.
- dP/dt is not an accurate measure because this increases with increasing preload and afterload.
- $(dP/dt)/P_{\text{ventricle}}$ is better. $P_{\text{ventricle}}$ is instantaneous ventricular pressure.
- Excess K^+ decreases contractility.
- Excess Ca^{++} causes spastic contraction, and low Ca^{++} causes cardiac dilation.


REGULATION OF STROKE VOLUME: AFTERLOAD




Measurement of Cardiac Output

- Electromagnetic flowmeter
- Indicator dilution (dye such as cardiogreen)
- Thermal dilution
- *Oxygen Fick Method*
- *$CO = (O_2 \text{ consumption} / (A-V O_2 \text{ difference}))$*

Electromagnetic flowmeter





$$q_1 = \dot{V}_O_2 \cdot C_{V_{O_2}}$$

q_2 = amount of Oxygen uptake by the lungs


$q_3 = \dot{V}_O_2 \cdot C_{A_{O_2}}$ and equals $= \dot{V}_O_2 \cdot C_{V_{O_2}} + O_2$ uptake

$$\text{Oxygen uptake} = \dot{V}_O_2 \{ C_{A_{O_2}} - C_{V_{O_2}} \}$$


$$\dot{V}_O_2 = \text{Oxygen uptake} / \{ C_{A_{O_2}} - C_{V_{O_2}} \}$$

Spirometer

A spirometer



Swan-Ganz catheter




O₂ Fick Problem

- If pulmonary vein O₂ content = 200 ml O₂/L blood
- Pulmonary artery O₂ content = 160 ml O₂ /L blood
- Lungs add 400 ml O₂ /min
- What is cardiac output?
- Answer: $400 / (200 - 160) = 10$ L/min

THE INDICATOR DILUTION PRINCIPLE

Indicator concentration



$$\text{Area} = \int_{t_2}^{t_1} dc \cdot dt$$

$$\text{Area} = \bar{C} * (t_2 - t_1)$$

(Rectangular)

$$\bar{C} = \text{Area} / (t_2 - t_1)$$


t_1

t_2

KMc

$$\text{Cardiac output} = \frac{q}{C} X \frac{60}{\text{duration in seconds}}$$

Thermodilution Method Curve




$$\text{AREA} = \int_{t1}^{t2} dT \cdot dt$$


VENOUS RETURN

- Definition: Volume of blood returns to either the left side or right side of the heart per minute
- $VR = CO = \Delta P/R$
- $VR = (\text{Venous pressure} - \text{Rt. Atrial pressure}) / \text{resistance to venous return}$

Effect of Venous Valves




Effect of Venous Valves





(a) Contracted skeletal muscles

(b) Relaxed skeletal muscles



Venous Valves



Effect Of Gravity on Venous Pressure




Vessel Structure and Function



Dilated and twisted appearance of varicose veins in the leg

Venous Pressure in the Body



- Compressional factors tend to cause resistance to flow in large peripheral veins.
- Increases in right atrial pressure causes blood to back up into the venous system thereby increasing venous pressures.
- Abdominal pressures tend to increase venous pressures in the legs.


Central Venous Pressure

- ❖ Pressure in the right atrium is called *central venous pressure*.
- ❖ *Right atrial pressure* is determined by the balance of the heart pumping blood out of the right atrium and flow of blood from the large veins into the right atrium.
- ❖ Central venous pressure is normally 0 mmHg, but can be as high as 20-30 mmHg.


Factors affecting Central Venous Pressure

⚙️ Right atrial pressure (RAP) is regulated by a balance between the ability of the heart to pump blood out of the atrium and the rate of blood flowing into the atrium from peripheral veins.


- ⚙️ Factors that increase RAP:
- ⚙️ -increased blood volume
 - ⚙️ -increased venous tone
 - ⚙️ - dilation of arterioles
 - ⚙️ -decreased cardiac function




Factors that Facilitate Venous Return



The Venous Return Curve





Venous Return (VR)


- Beriberi - thiamine deficiency \Rightarrow arteriolar dilatation $\Rightarrow \downarrow$ RVR
- (RVR= resistance to venous return)
because $VR = (MSFP - RAP) / RVR$
(good for positive RAP's)
- A-V fistula \Rightarrow (? RVR)
- \downarrow RVR
- C. Hyperthyroidism \Rightarrow (? RVR)
- \downarrow RVR

Venous Return (VR) (cont'd)

- Anemia \Rightarrow \downarrow RVR (why?)
- Sympathetics \Rightarrow MSFP
- Blood volume \Rightarrow MSFP + small \downarrow in RVR
- \downarrow Venous compliance (muscle contraction or venous constriction) \Rightarrow (? MSFP)
- MSFP

Factors Causing ↓ Venous Return

- ↓ Blood volume \Rightarrow ↓ MSFP
- ↓ Sympathetics \Rightarrow (? v. comp. and MSFP)
- Venous compliance and ↓ MSFP
- Obstruction of veins \Rightarrow (? RVR)
- RVR



Thank You

