

Doctor.021

no. 1

RS MICROBIOLOGY



Writer: 2020

Corrector: Noor Abu Hantash

Doctor: Microbiology department



the most common
microorganisms (bacteria and fungi) that may affect the
respiratory system.

Respiratory system

Microbiology laboratory section



-We will discuss common gram-positive bacteria that may cause throat infections. We will see how we can differentiate between the species.

-A throat swab culture is a test commonly used to diagnose bacterial infections in the throat

-staph and strep are among the commonest gram +ve bacteria

Staphylococcus

- Arranged in grape like clusters.
- Includes at least 40 species. The most common species associated with clinical infection are Staph aureus, Staph epidermidis, Staph hemolyticus, Staph hominins, and Staph albus.

Streptococcus

- Arranged in chains or diplococci.

THROAT SWAB

**Gram Positive
Coccus**



**Staphylococcus
Spp.**

**Streptococcus
Spp.**


GRAM STAIN



Gram+ve=blue to violet cocci

Staphylococcus

Streptococcus



Staph forms
grape-like
clusters



Strep
forms
pinpoint
chains

A- staphylococcus .albus:

Other staph species such as staph albus or epidermidis appear as white colonies on blood agar.

B- staphylococcus.Aureus

Staph aureus colonies appear yellow-golden and often present with hemolysis when grown on blood agar plates.

The golden appearance is the etymological root (origin) of the bacteria's name "aureus," as the word means golden in Latin.




Blood agar

Extra: The term "Staph albus" is an older name that was historically used for a group of staphylococcal bacteria, but it is not commonly used today. The modern nomenclature and classification have evolved, and Staphylococcus epidermidis is the accepted and more specific name for this particular bacterium.



Test for differentiation of Staphylococcus species

The major test reaction that should be used to differentiate between staphylococcus and streptococcus species is the catalase test




To differentiate between staph aureus and other staph species such as albicans, the mannitol salt agar (MSA) and coagulate tests should be used

Catalase test




- The catalase test is used to identify organisms that produce the enzyme catalase. This enzyme detoxifies hydrogen peroxide by breaking it down into water and oxygen gas. Bubbles resulting from the production of oxygen gas clearly indicates a catalase positive result




- staph species give +ve results
- strep species give -ve results

MSA


Mannitol salt agar media



Red → yellow (+ve mannitol fermenter)





Red → red
-ve



-The MSA is a selective and differential medium. Its high concentration of salt (7.5%) selects for members of the genus *Staphylococcus* since they can tolerate high saline levels. Organisms from other genera may grow but they grow weakly

-MSA also contains the sugar mannitol and the pH indicator phenol red. If an organism can ferment mannitol, an acidic byproduct is formed that will cause the phenol red in the agar to turn yellow. Staph aureus can ferment mannitol so its media will turn yellow while other Staph species such as epidermidis will not ferment mannitol and it will remain red in color.

Coagulase test



The coagulase test is used to differentiate staph aureus (+) from other staph species (-). Coagulase is an enzyme produced by staph aureus to convert soluble fibrinogen in plasma to insoluble fibrin.

-A suspension of the organism is suspended and incubated in plasma at 37C in a tube. Clot formation within four hours indicates a positive test, or the presence of staph aureus. Negative tubes should be left at room temperature overnight and re-examined the next day. This step is essential for some strains of staph aureus, including MRSA strains, as they produce a delayed clot which is rapidly lysed at 37C by staphylokinase

Streptococci are gram positive aerobic organisms that cause many disorders including pharyngitis, pneumonia, wound, skin infections, sepsis, and endocarditis. Three different types of strep are initially differentiated by their appearance when they are grown on sheep blood agar

-memorize the chart:



Streptococcus

α -hemolytic

green,
partial hemolysis

β -hemolytic

clear,
complete hemolysis

γ -hemolytic

no hemolysis

pneumoniae

optochin sensitive,
bile soluble,
capsule =>
quellung +

Viridans

mutans, sanguis
optochin resistant,
not bile soluble,
no capsule

pyogenes

Group A,
bacitracin sensitive

agalactiae

Group B,
bacitracin resistant

Enterococcus

E. faecalis,
E. faecium

Hemolysis on **sheep** blood agar

Blood Agar:

Shows three types of hemolysis


α Hemolysis

β Hemolysis


γ Hemolysis



Beta Hemolysis



Alpha Hemolysis




Gamma Hemolysis

Beta

Alpha

None



Differentiation between α -hemolytic streptococci

	Hemolysis	Optochin sensitivity
<i>S. pneumoniae</i>	α	Sensitive (≥ 14 mm)
<i>Viridans strep</i>	α	Resistant (≤ 13 mm)


Streptococcus viridians can be differentiated from *S. pneumoniae* using an optochin test.

Viridans streptococci are optochin resistant; they also lack either the polysaccharide-based capsule typical of *S. pneumoniae* or the Lancefield antigens of the pyogenic members of viridians




Optochin test

Streptococcus pneumoniae



Streptococcus pneumoniae strain on blood agar showing alpha hemolysis (green zone surrounding colonies). Note the **zone of inhibition** around a filter paper disc impregnated with optochin. (sensitive to optochin)

A zone of inhibition appears on a strep pneumoniae culture as it is sensitive to optochin



Streptococcus viridans strain on blood agar showing alpha hemolysis (green zone surrounding colonies). **No zone of growth inhibition (Resistant)** around a filter paper disc impregnated with optochin.

No zone of clearance


Streptococcus pneumoniae

Strep pneumoniae are gram positive, lancet shaped elongated cocci with a slightly pointed outer curvature. Usually they are seen in pairs (diplococci) but they can be singular or in short chains.



Streptococcus viridans

Strep viridans are gram positive (often elongated) cocci that form short to long chains.



Differentiation between β -hemolytic streptococci

	Hemolysis	Bacitracin sensitivity
<i>S. pyogenes</i>	β	Susceptible
<i>S. agalactiae</i>	β	Resistant

The bacitracin sensitivity test is used to distinguish group A *S. pyogenes* from other streptococci such as group B *S. agalactiae*.

When grown on blood agar, *S. pyogenes* is sensitive to bacitracin and will exhibit a zone of inhibition. While *S. agalactiae* will not be affected and will not have a zone of inhibition.



Strep agalactiae
There is no zone of
inhibition as it is
resistant to bacitracin.


Strep pyogenes
We can see the zone
inhibition indicating its
bacitracin sensitivity



Bacitracin test for *Streptococcus pyogenes*

Gamma hemolysis streptococcus

No, lack of hemolysis around the bacterium colony



**Enterococcus
Group D
- E.feacalis**



**Other than
Enterococcus
group D**

Non-enterococcus

The bile-esculin test is used to differentiate between enterococcus group D and non-enterococcus species from other strep viridans.

- Enterococcus species give a positive test.
- The bile-esculent test is based on the hydrolysis of esculin into glucose and esculetin by microorganisms that can produce esculinase enzyme.
- Esculetin then can interact with an iron salt, ferric citrate, in the medium to form a phenolic iron complex, which produces a dark brown or black color

Bile-Esculin



Diphtheroids


Diphtheroids are aerobic non-sporulating pleomorphic gram positive bacilli which are more uniformly stained than *Corynebacterium diphtheria*.

Gram Positive Cocco-bacilli

They lack metachromatic granules and are arranged as in what is known as the '**Chinese letter**' appearance. They are usually commensals in the skin and mucous membranes.

Arrangement as Chinese letter


文学家



Candida Species

Candidiasis is a fungal infection caused by a yeast known as Candida. Some species of Candida can cause infections in humans, most commonly *C. albicans*.


Candida normally lives on the skin and inside the body in places such as the mouth, throat, gut, and vagina, without causing any problems. In some circumstances, such as when taking a long course of Abx or a weakened immune system, the risk of Candida infection increases.




Candida Spp

- Larger than Bacteria
- Budding
 - Candida are unicellular fungi.
 - They may be spherical, elliptical or cylindrical shaped.
 - Their size varies greatly but they are generally larger than bacteria.
 - They typically grow asexually by budding.

Simple staining in lab



sabouraud dextrose agar




Sabouraud agar is agar growth media contains peptones. It is used to cultivate dermatophytes and other fungi (such as *Candida*) at 20C. It can also grow filamentous bacteria such as *Nocardia*. The pH of the media is adjusted to approximately 5.6 in order to enhance the growth of fungi, especially dermatophytes, and to slightly inhibit bacterial growth in clinical specimens. Yeast will grow as creamy white colonies while molds will grow as filamentous colonies with yeasty smell.



To Differentiate between C.albican and other Species

The germ tube test is a screening test used to differentiate *C. albicans* from other yeasts. When *Candida* is grown in human or sheep serum at 37C for three hours, it forms germ tubes. These germ tubes can be detected with wet KOH films as filamentous outgrowth extending from yeast cells. In this case, the sample is positive for *C. Albicans*



Germ tube





[Serum + candida]

You have to
memorize
colors^-^

Chrom agar

Chrom agar is a novel differential culture medium that facilitates the isolation and identification of some clinically important yeast species



***C. glabrata* : violet (dark pink)
glistening**




***C. albicans* : Green**

Chrom agar



C. Krusei isolates forms highly characteristic rough, spreading colonies with pale pink centers and white edges

**C. Krusei : rough
dry pale pink**




**C. Tropicalis
Blue grey**

Aspergillus Niger

Aspergillus niger is a fungus and one of the most common species of Aspergillus. It causes a disease called black mold on certain fruits and vegetables such as grapes, apricots, onions, and peanuts. It is a common contaminant of food. Aspergillus niger is one of the most common causes of otomycosis (fungal ear infection) which can cause pain or temporary hearing loss. In severe cases, it may damage the ear canal and tympanic membrane

Black colonies



Penicillium Spp.

Penicillium is a genus of ascomycetous fungi that is of major importance in the natural environment, in food spoilage, and in food and drug production. Some members of the genus produce penicillin, a molecule that is used as an antibiotic, which kills or stops the growth of certain kinds of bacteria. Penicillium species are occasional causes of infection in humans and the resultant disease is known generally as penicilliosis. Penicilliums have been isolated from patients with keratitis, endophthalmitis, otomycosis, necrotizing esophagitis, pneumonia, endocarditis, peritonitis and urinary tract infections.




Mycobacterium tuberculosis is a species of pathogenic bacteria in the family Mycobacteriaceae and is the causative agent of Tuberculosis.

It has an unusual waxy coating on its cell surface primarily due to the presence of mycolic acid. This coating makes the cell impervious to gram staining, and as a result M.

Tuberculosis can appear as either gram negative or gram positive.

So, acid fast stains, such as the Ziehl-Nielsen stain, are used to identify it with a microscope.



Lowenstein –Jensen Medium(LJ medium)

- Contain malachite green and egg albumin
- Media color : green
- Cell show :




Rough

Tough

Buff


Green → brown, granular colonies ("buff, rough and tough").

The LJ medium is a growth medium specially used for the culture of mycobacterium species, especially *M. Tuberculosis*. When grown on LJ medium, *M. Tuberculosis* appears as brown granular colonies sometimes called buff, rough, and tough colonies. The medium must be incubated for a significant length of time, usually 4 weeks, due to the slow proliferation time of this bacteria. The medium appears green, opaque and opalescent. The medium consists of malachite green, glycerol (which enhances the growth of *M. tuberculosis*), asparagine, potato starch, coagulated eggs, mineral salt solution, potassium dihydrogen phosphate, and magnesium sulfate.



Incubation Period = 4 weeks

Use glass bottles rather than petri dishes bcz of the long incubation time required, this type of bottles limits both chances of contamination and drying of culture media if the bottle is tightly closed



Put the media in covered tubes to avoid drying of media



ziehl neelsen= Acid fast stain



- Is used to stain TB species including MTB
- Mycobacterium Tuberculosis cell wall are waxed for that reason do heating while staining .
- Stain made of :




- **Carbol-fuchsin(primary stain)**

- Hydrochloric acid alcohol (3% HCL)
(de-colorizer)

- **Methylen blue(counter stain)**

When the smear is stained with carbol fuchsin, it solubilizes the lipoid always material present in the mycobacterial cell wall. With the application of heat, carbol fuchsin further penetrates through the lipoidal wall and enters into the cytoplasm. All cells now appear red. The smear is then decolorized with a decolorizing agent (3% HCl in 95% alcohol) but the acid fast cells are resistant due to the presence of large amounts of lipoidal material in their cell walls which prevents the penetration of decolorizing solution.


-Non-acid fast organisms lack the lipoidal material in their cell wall and are therefore easily decolorized, leaving colorless cells. Then, the smear is stained with a counter stain, methylene blue. Only decolorized cells absorb the counter stain and take its color and appear blue while acid-fast retain the red color



T.B Acid fast stain



Here, the M, tuberculosis cells appear red/pink in color while other cells in the background appear blue



1. Germ tube test is diagnostic for?

- A. Cryptococcus neoformans.
- B. Candida Tropicalis.
- C. Candida glabrata.
- D. Pseudomonas spp.
- E. None of the above.

Answer: E

2. Species is frequently associated with nosocomial infection with the following lab results:

- Gram-positive coccus arranged in chains.
- Beta hemolytic reaction in blood agar.
- Sensitive to bacitracin.

- A. Staphylococcus epidermidis.
- B. Streptococcus pyogenes (Group A).
- C. Enterococcus group D.
- D. Streptococcus pneumonia.
- E. Staphylococcus Albus.

Answer: B

3. Mannitol salt agar is a selective medium for?

- A. β hemolytic streptococci.
- B. Staphylococcus aureus.
- C. Corynebacterium diphtheria.
- D. Mycobacterium TB.
- E. Streptococcus group D.

Answer: B

4. This gram-positive coccobacillus micro-organism arrangement as Chinese letter is isolated from throat swab?

- A. Streptococcus viridans.
- B. Streptococcus group A.
- C. Streptococcus group B.
- D. Diphtheroid spp.
- E. None of the above.

Answer: D

1) **Staphylococcus aureus, which is WRONG?**

- A. Catalase +ve
- B. Cannot convert fibrinogen to fibrin
- C. Ferments mannitol salt
- D. Forms clusters in growth
- E. A+B

Answer: B

2) **Which of the following look green in chrome agar and form thick colonies in sabroud dextrose agar?**

- A. C. albicans (Ans)
- B. C. Tropicalis
- C. C. Krusie
- D. C. galbarta

Answer: A

3. **Which of the following is gram positive, catalase negative, positive bile- esculin?**

- A. E. faecalis
- B. Strep viridans

Answer: A

4. The correct order in zeihl Nelsen staining is:

- A. Carbon fuchsin, heating, acid-alcohol, methylene blue
- B. Carbon fuchsin, acid-alcohol, methylene blue, heating
- B. Carbon fuchsin, acid-alcohol, methylene blue, heating
- C. Methylene blue, heating, acid-alcohol, carbon fuchsin
- D. None of the above

Answer: A

5. Streptococcus pneumoniae, one is incorrect:

- A. Diplococci.
- B. have capsules.
- C. lysed by bile.
- D. Resistant to optochin.
- E. Produce a hemolysis.

Answer: D

6. The microorganism which is catalase Negative and sensitive to Optochin is:

- A. Beta hemolytic streptococcus group A.
- B. Streptococcus Pneumoniae.
- C. Enterococcus group D.
- D. Staphylococcus aureus.
- E. Neisseria Spp.

Answer: B

7. Which of the following is an appropriate media for all fungi?

- A. SDA.
- B. Chrome Agar.

Answer: A

8. Which of the following is sensitive to bacitracin?


- A. Enterococcus.
- B. Strep agalactiae.
- C. Strep pneumonia.
- D. Strep pyogenes.

Answer: D

9. The following media is:

- A. Chrome agar.
- B. Lowenstein-Jensen.
- C. Sabouraud dextrose agar.

Answer: A



16. The type of fungus that produces the Blue color on chrom agar media:

- A. Candida tropicalis.
- B. Candida glabrata.
- C. None of the mentioned.
- D. Candida krusei.
- E. Candida albican.

Answer: A

17. The microorganism which is catalase Negative and sensitive to Optochin is:

- A. Beta hemolytic streptococcus group A.
- B. Streptococcus Pneumoniae.
- C. Enterococcus group D.
- D. Staphylococcus aureus.
- E. Neisseria Spp.

Answer: B

18. The major test reaction should be used to differentiate between staphylococcus and streptococcus species?

Answer: Catalase test

19. Which of the following test is used to differentiate between staph aureus and other staph species?

Answer: mannitol salt agar (MSA) and coagulate tests

20. Which of the following test is used to differentiate between enterococcus group D and non-enterococcus species?


Answer: Bile-esculin test

Collected by: Shahed Atiyat

10. The test shows:

- A. Strep. Pneumonia.
- B. Strep. Viridans.
- C. Strep. Pyogenes.
- D. Strep. Agalactiae.

Answer: B



optochin test

11. Which of the following mediums used for the culture of M. Tuberculosis?

Answer: Lowenstein –Jensen Medium

12. The organism with alpha hemolysis and it is optochin resistant?

Answer: Viridans streptococci

13. Which of the following is sensitive to bacitracin?

- A. Enterococcus.
- B. Strep agalactiae.
- C. Strep pneumonia.
- D. Strep pyogenes.

Answer: D



V2: check the highlight as follows:

Identification of Streptococci

CATALASE TEST

↓ - ●●●●

Streptococcaceae

↓

Distinguishing tests

	<u>Disks</u>		<u>Ability to Grow in:</u>	
	<u>Optochin</u>	<u>Bacitracin</u>	<u>6.5% NaCl</u>	<u>Bile esculin</u>
<i>S. pneumoniae</i>	S	R	-	-
<i>S. pyogenes</i>	R	S	-	-
<i>E. faecalis</i>	R	R	+	+
Nonenterococcal Gp D	R	R	-	+
Viridans streptococci	R	R	-	-

