

Urinary System: Renal Physiology for Medical Students, L11-12



Chapter 30: Acid-Base Regulation in the Kidney

Reference: Guyton & Hall, Jordanian first edition


Dr. Ebaa M. Alzayadneh, PhD.

Email: e.zayadneh@ju.edu.jo


2024

Objectives

- Identify the mechanisms by which the kidney can maintain Acid-Base Balance
- Identify the most important buffers in different body fluids
- Identify the cellular mechanisms of for HCO_3^- reabsorption and $\text{Na}^+ - \text{H}^+$ exchange or H^+ secretion in the nephron
- Understand the mechanism of renal compensations for Acid-Base Disorders.
- To be able to determine the type of acid-base imbalance from given lab results and figure out the compensatory changes.
- To know how to use “Anion Gap” as a Diagnostic Tool for Metabolic Acidosis
- Identify main clinical conditions that are accompanied with acid-base imbalance



Audio-Visual Aid



- Please watch this video to help introducing you to the topic of this lecture

[Acid Base Balance, Animation. - YouTube](#)



Search

Pulmonary regulation

$$\text{CO}_2 + \text{H}_2\text{O} \leftrightarrow \text{H}_2\text{CO}_3 \leftrightarrow \text{H}^+ + \text{HCO}_3^-$$

© ALILA MEDICAL MEDIA
License this video at
www.AlilaMedicalMedia.com

A decrease in pH is sensed by central or arterial chemoreceptors and leads to deeper, faster
(Communicating vessels)

Acid Base Balance, Animation.



Mechanisms of Hydrogen Ion Regulation



[H⁺] is precisely regulated at $3-5 \times 10^{-8}$ moles/L
(pH range 7.2 -7.4)

1. Body fluid chemical buffers (rapid but temporary)

- bicarbonate
- ammonia
- proteins
- phosphate

2. Lungs (rapid, eliminates CO₂)

↑[H⁺] → ↑ ventilation → ↑ CO₂ loss

3. Kidneys (slow, powerful); eliminates non-volatile acids

- secretes H⁺
- reabsorbs HCO₃⁻
- generates new HCO₃⁻

Buffer Systems in the Body

Bicarbonate : most important ECF buffer



Phosphate : important renal tubular buffer



Ammonia : important renal tubular buffer



Proteins : important intracellular buffers



(60-70% of buffering is in the cells)

Importance of Buffer Systems

Normal H^+ concentration = 0.00004 mmol/L


Amount of non-volatile acid produced
~ 60-80 mmol/day

80 mmol /42 L = 1.9 mmol/L

= 47,500 times > normal H^+ concentration

PH ----6.8-8 lives for hours

Bicarbonate Buffer System



$$\text{pH} = \text{pK} + \log \frac{\text{HCO}_3^-}{\alpha \text{ pCO}_2} \quad \begin{array}{l} \alpha = 0.03 \\ \text{pK} = 6.1 \end{array}$$

Effectiveness of buffer system depends on:

- concentration of reactants
- pK of system and pH of body fluids

Titration curve for bicarbonate buffer system.





Figure 30-1.

Bicarbonate Buffer System

Is the most important buffer in extracellular fluid even though the concentration of the components are low and pK of the system is 6.1, which is not very close to normal extracellular fluid pH (7.4).

Reason: the components of the system (CO_2 and HCO_3^-) are closely regulated by the lungs and the kidneys

Respiratory Regulation of Acid-Base Balance



Feedback Gain = 1.0 to 3.0
(corrects 50 to 75 %)

Renal Regulation of Acid-Base Balance

- Kidneys eliminate non-volatile acids (H_2SO_4 , H_3PO_4) (~ 80 mmol/day)
- Filtration of HCO_3^- (~ 4320 mmol/day)
- Secretion of H^+ (~ 4400 mmol/day)
- Reabsorption of HCO_3^- (~ 4319 mmol/day)
- Production of new HCO_3^- (~ 80 mmol/day)
- Excretion of HCO_3^- (1 mmol/day)

Kidneys conserve HCO_3^- and excrete acidic or basic urine depending on body needs

Reabsorption of bicarbonate (and H⁺ secretion) in different segments of renal tubule.

Key point:
For each HCO₃⁻ reabsorbed, there must be a H⁺ secreted




Figure 30-4.

Mechanisms for HCO_3^- reabsorption and $\text{Na}^+ - \text{H}^+$ exchange in proximal tubule and thick loop of Henle





Figure 30-5.

HCO_3^- reabsorption and H^+ secretion in intercalated cells of late distal and collecting tubules



Minimal
pH ~4.5


Figure 30-6.

Renal Regulation of Acid-Base Balance

- Kidneys eliminate non-volatile acids (H_2SO_4 , H_3PO_4) (~ 80 mmol/day)
- Filtration of HCO_3^- (~ 4320 mmol/day)
- Secretion of H^+ (~ 4400 mmol/day)
- Reabsorption of HCO_3^- (~ 4319 mmol/day)
- Production of new HCO_3^- (~ 80 mmol/day)
- Excretion of HCO_3^- (1 mmol/day)

Kidneys conserve HCO_3^- and excrete acidic or basic urine depending on body needs

Regulation of H⁺ secretion




$$\text{pH} = \text{pK} + \log \frac{[\text{HCO}_3^-]}{\alpha \text{ pCO}_2}$$

- Increased pCO₂ increases H⁺ secretion
i.e. respiratory acidosis
- Increased extracellular H⁺ increases H⁺ secretion
i.e. metabolic or respiratory acidosis
- Increased tubular fluid buffers increases H⁺ secretion
i.e. metabolic or respiratory acidosis

Renal Compensations for Acid-Base Disorders

- Acidosis:
 - increased H^+ secretion
 - increased HCO_3^- reabsorption
 - production of new HCO_3^-
- Alkalosis:
 - decreased H^+ secretion
 - decreased HCO_3^- reabsorption
 - loss of HCO_3^- in urine

In acidosis all HCO_3^- is titrated and excess H^+ in tubule is buffered



Importance of Renal Tubular Buffers

$$\begin{aligned}\text{Minimum urine pH} &= 4.5 \\ &= 10^{-4.5} \\ &= 3 \times 10^{-5} \text{ moles/L}\end{aligned}$$

i.e. the maximal $[\text{H}^+]$ of urine is 0.03 mmol/L

Yet, the kidneys must excrete, under normal conditions, at least 60 mmol non-volatile acids each day. To excrete this as free H^+ would require :

$$\frac{60 \text{ mmol}}{.03 \text{ mmol/L}} = 2000 \text{ L per day !!!}$$

Buffering of secreted H^+ by filtered phosphate ($NaHPO_4^-$) and generation of “new” HCO_3^-




Figure 30-7.

Phosphate as a Tubular Fluid Buffer


There is a high concentration of phosphate in the tubular fluid; $pK = 6.8$

Phosphate normally buffers about 30 mmol/day H^+ (about 100 mmol/day phosphate is filtered but 70 % is reabsorbed)

Phosphate buffering capacity does not change much with acid-base disturbances (phosphate is not the major tubular buffer in chronic acidosis)



Phosphate and Ammonium Buffering In Chronic Acidosis



Production and secretion of NH_4^+ and HCO_3^- by proximal, thick loop of Henle, and distal tubules




Figure 30-8.

Buffering of hydrogen ion secretion by ammonia (NH_3) in the collecting tubules.




Figure 30-9.

Quantification of Normal Renal Acid-Base Regulation

Total H⁺ secretion

= 4320 mEq of H⁺ secreted (HCO₃) + 60 mEq of H⁺ non-volatile = 4380

Total H⁺ secretion = 4380 mmol/day

= HCO₃⁻ reabsorption (4320 mmol/d)

+ titratable acid (NaHPO₄⁻) (30 mmol/d)

+ NH₄⁺ excretion (30 mmol/d)

Net H⁺ excretion =

H⁺ excreted by buffers not bicarbonate (= new bicarb) - new H⁺ added to blood (= HCO₃⁻ excreted)

Net H⁺ excretion = 59 mmol/day

= titratable acid (30 mmol/d)

+ NH₄⁺ excretion (30 mmol/d)

- HCO₃⁻ excretion (1 mmol/d) (or new H to blood)

Normal Renal Acid-Base Regulation

Net addition of HCO_3^- to body
(i.e. net loss of H^+)

$$\begin{array}{rcl} \text{Titratable acid} & & = 30 \text{ mmol/day} \\ + \text{NH}_4^+ \text{ excretion} & = & 30 \text{ mmol/day} \\ - \text{HCO}_3^- \text{ excretion} & = & 1 \text{ mmol/day} \\ \text{Total} & & = 59 \\ \text{mmol/day} & & \end{array}$$

Renal Compensation for Acidosis

Increased addition of HCO_3^- to body by kidneys
(increased H^+ loss by kidneys)

Titratable acid	= 35 mmol/day (small increase)
NH_4^+ excretion	= 165 mmol/day (increased)
HCO_3^- excretion	= 0 mmol/day (decreased)
Total	= 200 mmol/day

This can increase to as high as 500 mmol/day


Renal Compensation for Alkalosis

Net loss of HCO_3^- from body
(i.e. decreased H^+ loss by kidneys)

Titratable acid	=	0 mmol/day (decreased)
NH_4^+ excretion	=	0 mmol/day (decreased)
HCO_3^- excretion	=	80 mmol/day (increased)
Total	=	80 mmol/day

HCO_3^- excretion can increase markedly in alkalosis

Classification of Acid-Base Disorders from plasma pH, pCO₂, and HCO₃⁻



$$\text{pH} = \text{pK} + \log \frac{\text{HCO}_3^-}{\alpha \text{ pCO}_2}$$

Acidosis : pH < 7.4

- metabolic : ↓ HCO₃⁻
- respiratory : ↑ pCO₂

Alkalosis : pH > 7.4

- metabolic : ↑ HCO₃⁻
- respiratory : ↓ pCO₂


Renal Compensations for Acid-Base Disorders

- Acidosis:
 - increased H^+ excretion
 - increased HCO_3^- reabsorption
 - production of new HCO_3^-
- Alkalosis:
 - decreased H^+ excretion
 - decreased HCO_3^- reabsorption
 - loss of HCO_3^- in urine

Renal Responses to Respiratory Acidosis




Respiratory acidosis : \downarrow pH \uparrow pCO₂ \uparrow HCO₃⁻




Renal Responses to Metabolic Acidosis

Metabolic acidosis : \downarrow pH \downarrow pCO₂ \downarrow HCO₃⁻




Renal Responses to Respiratory Alkalosis

Respiratory alkalosis : \uparrow pH \downarrow pCO₂ \downarrow HCO₃⁻



Renal Responses to Metabolic Alkalosis

Metabolic alkalosis : \uparrow pH \uparrow pCO₂ \uparrow HCO₃⁻





Question

The following data were taken from a patient:

urine volume = 1.0 liter/day

urine HCO_3^- concentration = 2 mmol/liter

urine NH_4^+ concentration = 15 mmol/liter

urine titratable acid = 10 mmol/liter

- What is the daily net acid excretion in this patient ?
- What is the daily net rate of HCO_3^- addition to the extracellular fluids ?



Question

The following data were taken from a patient:

urine volume = 1.0 liter/day

urine HCO_3^- concentration = 2 mmol/liter


urine NH_4^+ concentration = 15 mmol/liter

urine titratable acid = 10 mmol/liter

$$\begin{aligned}\text{net acid excretion} &= \text{Titr. Acid} + \text{NH}_4^+ \text{ excret} - \text{HCO}_3^- \\ &= (10 \times 1) + (15 \times 1) - (1 \times 2) \\ &= 23 \text{ mmol/day}\end{aligned}$$

$$\text{net rate of } \text{HCO}_3^- \text{ addition to body} = 23 \text{ mmol/day}$$

Classification of Acid-Base Disorders from plasma pH, pCO₂, and HCO₃⁻



$$\text{pH} = \text{pK} + \log \frac{\text{HCO}_3^-}{\alpha \text{ pCO}_2}$$

Acidosis : pH < 7.4

- metabolic: ↓ HCO₃⁻
- respiratory: ↑ pCO₂


Alkalosis : pH > 7.4

- metabolic: ↑ HCO₃⁻
- respiratory: ↓ pCO₂



Classification of Acid-Base Disturbances

Disturbance	Plasma			Compensation
	pH	HCO ₃ ⁻	pCO ₂	
metabolic acidosis	↓	↓	↓	↑ ventilation ↑ renal HCO ₃ production
respiratory acidosis	↓	↑	↑	↑ renal HCO ₃ production
metabolic alkalosis	↑	↑	↑	↓ ventilation ↑ renal HCO ₃ excretion
respiratory alkalosis	↑	↓	↓	↑ renal HCO ₃ excretion



	pH	HCO ₃ ⁻	CO ₂
Metabolic acidosis	↓	↓	Normal
Metabolic alkalosis	↑	↑	Normal
Metabolic acidosis with respiratory compensation	↓	↓	↓
Metabolic alkalosis with respiratory compensation	↑	↑	↑

Test	Normal	Decrease Value	Increase Value
pH	7.35-7.45	Acidosis	Alkalosis
PaCO ₂	35-45	Alkalosis	Acidosis
HCO ₃	22-26	Acidosis	Alkalosis
PaO ₂	80-100	Hypoxemia	O ₂ therapy
SaO ₂	95-100%	Hypoxemia	----- -





Question

A plasma sample revealed the following values
in a patient: norm for PCO_2 35-45, HCO_3^- 22-26

$$\text{pH} = 7.12$$

$$\text{PCO}_2 = 50$$

$$\text{HCO}_3^- = 18$$

diagnose this patient's acid-base status :

acidotic or alkalotic ?

Acidotic

respiratory, metabolic, or both ?

Both

Mixed acidosis: metabolic and respiratory acidosis

Mixed Acid-Base Disturbances

Two or more underlying causes of acid-base disorder.

pH= 7.60

pCO₂ = 30 mmHg

plasma HCO₃⁻ = 29 mmol/L

What is the diagnosis?

Mixed Alkalosis

- Metabolic alkalosis : increased HCO₃⁻
- Respiratory alkalosis : decreased pCO₂



Question

A patient presents in the emergency room and the following data are obtained from the clinical labs:


plasma pH= 7.15, $\text{HCO}_3^- = 8 \text{ mmol/L}$, $\text{pCO}_2 = 24 \text{ mmHg}$

This patient is in a state of:

1. metabolic alkalosis with partial respiratory compensation
2. respiratory alkalosis with partial renal compensation
3. metabolic acidosis with partial respiratory compensation
4. respiratory acidosis with partial renal compensation

Acid-Base Disturbances

- Metabolic Acidosis : $\downarrow \text{HCO}_3^- / \text{pCO}_2$ in plasma
 \downarrow ($\downarrow \text{pH}, \text{HCO}_3^-$)
 - aspirin poisoning ($\uparrow \text{H}^+$ intake)
 - diabetes mellitus ($\uparrow \text{H}^+$ production)
 - diarrhea (HCO_3^- loss)
 - renal tubular acidosis ($\downarrow \text{H}^+$ secretion, $\downarrow \text{HCO}_3^-$ reabs.)
 - carbonic anhydrase inhibitors ($\downarrow \text{H}^+$ secretion)
-





Anion Gap as a Diagnostic Tool

In body fluids: total cations = total anions

Cations (mEq/L)

Na⁺ (142)

K⁺ (4)

Ca⁺⁺ (5)

Mg⁺⁺ (2)

Total (153)

Anions (mEq/L)

Cl⁻ (108)

HCO₃⁻ (24)

Unmeasured

Proteins (17)

Phosphate,

Sulfate,

lactate, etc (4)

(153)

Anion Gap as a Diagnostic Tool

In body fluids: total cations = total anions

$$\text{Na}^+ = \text{Cl}^- + \text{HCO}_3^- + \text{unmeasured anions}$$

$$\text{unmeasured anions} = \text{Na}^+ - \text{Cl}^- - \text{HCO}_3^- = \text{anion gap}$$

$$= 142 - 108 - 24 = 10 \text{ mEq/L}$$

$$\text{Normal anion gap} = 8 - 16 \text{ mEq / L}$$

Anion Gap in Metabolic Acidosis

- loss of HCO_3^- = normal anion gap

↔ anion gap = Na^{\uparrow} - Cl^{\downarrow} - HCO_3^-
hyperchloremic metabolic acidosis

- \uparrow unmeasured anions = \uparrow anion gap

\uparrow anion gap = $\text{Na}^{\leftrightarrow}$ - Cl^{\downarrow} - HCO_3^-
normochloremic metabolic acidosis
i.e. diabetic ketoacidosis, lactic acidosis,
salicylic acid, etc.



Use of “Anion Gap” as a Diagnostic Tool for Metabolic Acidosis

Increased Anion Gap (normal Cl^-)

- diabetes mellitus (ketoacidosis)
- lactic acidosis
- aspirin (acetylsalicylic acid) poisoning
- methanol poisoning
- starvation

Normal Anion Gap (increased Cl^- , hyperchloremia)

- diarrhea
- renal tubular acidosis
- Addison’ disease
- carbonic anhydrase inhibitors



Question

Laboratory values for an uncontrolled diabetic patient include the following:

arterial pH = 7.25

Plasma $\text{HCO}_3^- = 12$

Plasma $\text{P}_{\text{CO}_2} = 28$

Plasma $\text{Cl}^- = 102$

Plasma $\text{Na}^+ = 142$

Metabolic Acidosis

Respiratory Compensation

What type of acid-base disorder does this patient have?

What is his anion gap ?

$$\text{Anion gap} = 142 - 102 - 12 = 28$$



Question


Which of the following are the most likely causes of his acid-base disorder?

- a. diarrhea
- b. diabetes mellitus**
- c. Renal tubular acidosis
- d. primary aldosteronism

Acid-Base Disturbances


- Respiratory Acidosis : ↓ HCO_3^- / pCO_2 in plasma
↓ (↑ pH, pCO_2)


- brain damage
- pneumonia
- emphysema
- other lung disorders




Acid-Base Disturbances

- Metabolic Alkalosis : $\uparrow \text{HCO}_3^- / \text{pCO}_2$ in plasma
 \uparrow ($\uparrow \text{pH}$, HCO_3^-)
 - increased base intake (e.g. NaHCO_3)
 - vomiting gastric acid
 - mineralocorticoid excess
 - overuse of diuretics (except carbonic anhydrase inhibitors)







Acid-Base Disturbances

• Respiratory Alkalosis : $\uparrow \text{HCO}_3^- / \text{pCO}_2$ in plasma
 \uparrow ($\downarrow \text{pH}$, pCO_2)

- high altitude

- psychic (fear, pain, etc)



$$\uparrow \text{pH} = \text{pK} + \log \frac{\text{HCO}_3^-}{\alpha \text{pCO}_2}$$



Question

Laboratory values for a patient include the following:

arterial pH = 7.34

Plasma $\text{HCO}_3^- = 15$

Plasma $\text{P}_{\text{CO}_2} = 29$

Plasma $\text{Cl}^- = 118$

Plasma $\text{Na}^+ = 142$

Metabolic Acidosis

Respiratory Compensation

What type of acid-base disorder does this patient have?

What is his anion gap ?

$$\text{Anion gap} = 142 - 118 - 15 = 9 \quad (\text{normal})$$



Question

Which of the following are the most likely causes of his acid-base disorder?

a. diarrhea

b. diabetes mellitus

c. aspirin poisoning

d. primary aldosteronism



Indicate the Acid -Base Disorders in Each of the Following Patients

pH	HCO ₃ ⁻	PCO ₂	Acid-Base Disorder ?
7.34	15	29	Metabolic acidosis
7.49	35	48	Metabolic alkalosis
7.34	31	60	Respiratory acidosis
7.62	20	20	Respiratory alkalosis
7.09	15	50	Acidosis: respiratory + metabolic