



ISCHEMIC HEART DISEASE

Prof Akram Saleh, MD,FRCP
Consultant Invasive Cardiologist

Edited by: Ruaa Adeb

A 60 YEAR OLD MALE, SMOKER, DM PRESENTED C/O CHEST PAIN OF 6 MONTHS DURATION

Retrosternal

Heavy

PPT by exertion → exacerbated

Relieved by rest

Last about 5 minutes

DIAGNOSIS: *stable angina*

Ischemic Heart Disease (IHD)

chronic coronary syndrome
acute coronary syndrome

Basic: coronary circulation

Myocardial oxygen supply

What is IHD

Causes of IHD

Manifestations of IHD

Treatment

Heart Anatomy

In the middle mediastinum

- The heart is about the size of a fist and weighs 300-450 gm
- The average beat per minute is 70
- The average adult heart pumps about 6000-7500 liters of blood per day.

↳ Reciprocal changes: ST depressions on the other leads than the MI leads


↳ Tombstone ST elevations are very severe and high mortality

↳ ST elevation \curvearrowright convex \cap → classic for MI
 \curvearrowleft concave \cup → classic for pericarditis

Coronary Anatomy

↳ 1st branch of aorta

there are some collaterals (not functioning)
to function need → transient ischemia



Lateral
I, AVL, ^{LCX}
V5-V6

→ LAD

Anterior /
Septal
V1-V4 ^{LAD}

Inferior
II, III, aVF

PDA

↳ Right heart dominance
يعني غالباً الـ RA يكون
باد right

Coronary Circulation physiology

at rest

1- Flow during basal cardiac circulation: 70-80 ml/min/100gm

2- Flow during maximal cardiac work: 300-400ml/min/100gm

3- High oxygen extraction: 65%-75% (**fixed**)

need ↑ O₂

need ↑ coronary flow

↳ if not → ischemia

4- 80% of coronary flow occurs in diastole

5- Collateral pathways is anatomically present but not functioning


6- Cardiac tissue hypoxia is the potent stimulus to open the collateral

Ischemic Heart Disease

Myocardial O₂
demand

→ to avoid tachycardia: give β -blocker if no contraindication like asthma

- 1- Heart rate
- 2- ↑ Contractility → ↑ O₂ demand
- 3- Wall tension
- 4- Muscle mass (wall thickness)



Myocardial O₂
supply

1- Coronary flow (patency of coronary artery)

2- Hemoglobuline level ↳ healthy or not

Fixed ↳ 3- Myocardial oxygen extraction

Lung disease ↳ 4- Arterial oxygen saturation

CAUSES OF Myocardial ischemia

Reduced Myocardial O₂ Supply

1-Coronary artery disease

(atherosclerosis and non-atherosclerosis)

2-decrease flow of oxygenated blood:

Sever Anemia
carboxyhemoglobinemia
Hypotension

Increased Myocardial O₂ Demand

1-Left Ventricular Hypertrophy:

cause angina due to ↑ in myocardial O₂ demand {
hypertension
aortic stenosis
hypertrophic cardiomyopathy

2- Increase cardiac output:

Thyrotoxicosis
Rapid Tachyarrhythmias

Causes of coronary artery disease

5%

Atherosclerosis 95%

Risk factors

Nonatherosclerosis

1-Arteritis like: vasculitis
(SLE, RA, Takayasu ,..)

2-Embolism

3-Coronary mural thickening
(amyloidosis, radiation
therapy,..)

4-Coronary luminal
narrowing: coronary spasm,
aortic dissection

5-Congenital coronary
artery anomalies

Risk Factors for Cardiovascular Disease

Modifiable

4 major risk factors

- Hypertension
- Smoking → passive & active
- Hyperlipidaemia
 - Raised LDL-C
 - Low HDL-C
 - Raised triglycerides
- Diabetes mellitus
- Dietary factors
- Lack of exercise
- Obesity → trunkal
- Homocysteinemia
- Lipoprotein a
- Gout
- Thrombogenic factors: fibrinogen, factors V, VII
- Excess alcohol consumption



Non-modifiable

- Personal history of CVD
- Family history → 1st degree of CVD
M < 55 F < 65
- Age: M > 45, F > 55
- Gender M > F (Premenopausal)
- Personality type A
- Genetic factors: ACE gene

Prevalence of Modifiable Conventional Risk Factors in Patients With CAD in Jordan

At least one RF
Diabetes
HTN
Current smoking
C>240
TG>150
HDL<40

1534	
Men 1202 (78%)	Women 332 (22%)





95% of Patients With CAD in Jordan Have at least one of the Modifiable Conventional Risk Factors

Number of conventional risk factors in individuals with CAD in Jordan

Men with CAD

Diabetes
HTN
Smoking
C>240
TG>150
HDL<40



Pathogenesis of Atherosclerotic Plaques

(mechanical shear stresses, biochemical, immunological, inflammation, genetics abnormalities)

Endothelial damage (Dysfunction)



Protective response results in production of cellular adhesion molecules

(Cytokines, Chemokines, Growth factors)



Monocytes and T lymphocytes attach to 'sticky' surface of endothelial cells



Migrate through arterial wall to subendothelial space



Macrophages take up oxidised LDL-C




Lipid-rich foam cells




Fatty streak and plaque

Fatty Streak Formation in Atherosclerosis




Formation of the Complicated Atherosclerotic Plaque




Characteristics of Unstable and Stable Plaque

Unstable



Stable



- too late presentation
- start Tx before symptoms

« tip of the iceberg »

Cardiac Event


Clinical Test



Asymptomatic

Myocardial Ischemia: Chest pain is the tip of the iceberg

Major Manifestations of Atherothrombosis



Cerebrovascular disease

Coronary artery disease

Renal artery stenosis

Visceral arterial disease

Peripheral arterial disease (PAD)

The Spectrum of presentations

Myocardial Ischemia

**Stable
Angina**

**Unstable
Angina**

**Non-ST
Elevated MI
(NSTEMI)**

**ST
Elevated MI
(STEMI)**

**Sudden
Death**



Acute Coronary Syndromes

Thrombus present in the artery

Clinical Manifestations of Atherosclerosis

Coronary heart disease

- *Asymptomatic, Angina pectoris, variant angina, unstable angina, myocardial infarction, congestive heart failure (CHF), arrhythmias, and sudden cardiac death.*



Cerebrovascular disease

- *Transient ischaemic attack, stroke*

Peripheral vascular disease

- *Intermittent claudication, gangrene, cold feet, painful feet, impotence*

IHD-clinicopathological correlation

1- stable angina: stenosis > 70% luminal narrowing

2-variant angina: increase coronary tone
30% normal coronaries

3-unstable angina: rupture plaque
subocclusive thrombus (incomplete occlusion)
progress to myocardial infarction 15-30%

4-myocardial infarction: rupture plaque
occlusive thrombus (complete occlusion)

Stable angina

Commonest form of angina

Causes: imbalance between demand and supply

Symptom: chest pain

Location: central chest (others)

Radiation: arm(s), neck, jaw

Character : squeezing, pressure, heaviness,..

Duration: 2-10 minutes

Precipitating factors: exertion, emotional upset, heavy meal, sexual intercourse, cold weather

Relieving factors: nitrate, rest

Associated symptoms: dyspnea, diaphoresis, nausea

Classes of angina: 1- 4


Physical Examination: normal, sign of risk factors, peripheral vascular disease

Symptoms of Angina



Angina can spread anywhere between the belly button and the jaw, including to the shoulder, arm, elbow or hand- usually on the left side.

typically : chest pain



Angina → clinical diagnosis

ECG may be normal

Angina Chest Pain: Clinical Diagnosis



CAUSES OF ANGINA

Reduced Myocardial O₂ Supply

1-Coronary artery disease

2-Sever Anemia < 9 gm/dl

Increased Myocardial O₂ Demand

1-Left Ventricular Hypertrophy:

hypertension

aortic stenosis

hypertrophic cardiomyopathy

2- Rapid Tachyarrhythmias

NYHA Grading of cardiac symptoms

same classes of dyspnea (angina / dyspnea)

Grade 1:

Cardiac disease without resulting limitation of physical activity.

ordinary Ordinary physical activity does not cause chest pain (dyspnea).

Grade 2:

ordinary Slight limitation of physical activity. Comfortable at rest. Ordinary physical activity result in chest pain (dyspnea).

Grade 3:

ordinary moderate limitation in physical activity. Comfortable at rest. Less than ordinary activity causes symptoms

Grade 4:

Rest sever limitation: symptoms at rest. *unstable*

normal ECG will not rule out
any cardiac disease

→ no specific signs, normal exam

Stable angina- Diagnosis

↳ auscultate coronary for bruits


- **History : angina pectoris is clinical diagnosis**
- **Physical exam** *normal*
- **Electrocardiogram: 12 ECG, 24 ECG**
- 60-70% specific ← ▪ **Stress ECG : diagnostic and prognostic information**
- **Radioactive studies: thalium scan,..**
- **Echocardiography**
- **CT Coronary angiography**
- **Serum lipid(LDL, HDL, TG), FBG,CBC** → *anemia*
- **Coronary angiography**


Types of stress test

- *Exercise tolerance test: ST segment depression*
- *Exercise or dobutamine Echocardiogram: Wall motion abnormalities*
- *Exercise or dipyridamole Thallium: Decrease uptake of the nuclear isotope during exercise*

ST depression
subendocardial
ischemia

Resting Electrocardiogram






Types of stress test

- *Exercise or dipyridamole Thallium: Decrease uptake of the nuclear isotope during exercise*


Normal Myocardial Perfusion

Reversible ischemia = stable angina



Infarct = constant ischemia


Myocardial Ischemia



ischemic → hyperkinetic





الطبيعي، إنفاكوه مشدودة و
ما بتتحرك

Myocardial Infarction



Types of stress test

- *Exercise or dobutamine Echocardiogram: Wall motion abnormalities*



Imaging Techniques Used to Assess Atherosclerosis

Invasive techniques



- *Coronary angiography*
- *Intravascular ultrasound (IVUS)*

Non-invasive techniques

- *Magnetic resonance imaging (MRI)*
- *Computed tomography (CT)*
- *Ultrasound (B-mode)*


Intravascular Ultrasound (IVUS) Showing Atheromatous Plaque

Angiogram



IVUS

normal vessel



atheroma

HUSAM TAWFEEQ ESREWY

ID: 1032893

* 05/10/1974

Study 1

05/10/2017

12:22:06


1 IMA 56 FRM 1 AFPS 16

H

Jordan University Hospital

AXIOM-Artis

HFS



R

Coro 3040H.HT

Coro 3040H.HT

SINGLE PLANE\SINGLE A

CAU 34

RAO 5


W: 153

C: 125



Cardiac CATH





CARDIAC CT ANGIOGRAPHY

Management goals of stable angina

Main goals:

- **To improve prognosis (mortality reduction)**
 - *Modification of risk factors*
 - *Aspirin*
 - *Lipid-lowering therapy*
 - *ACE-Inhibitor*
 - *Revascularization procedures (PTCA, CABG)*
- **To decrease anginal symptoms**
 - *Medical treatment*

Treatment of stable angina

1- General measures

**2- Medical therapy: Increase O₂ supply
Decrease O₂ demand**

**3-Revascularization: PCI (percutaneous coronary intervention)
CABG (coronary artery bypass grafting)**

TREATMENT OF STABLE ANGINA

General Measures

- **Correction of established risk factors(reversible)**
- **weight reduction (ideal body weight)**
- **Aerobic exercise:**
improve functional capacity, well-being sensation
- **Treatment of: anemia, thyrotoxicosis, arrhythmias,..**

MEDICAL THERAPY OF STABLE ANGINA

Prognostic: **Aspirin, Statines, ACEI**

anti-platelet

β -B \rightarrow \downarrow HR
 \uparrow time for flow
 \uparrow diastole time

Symptomatic: **Nitrate, B-, CA-blocker, (nicorandil, ranolazine, ivabradine)**

CAB

INCREASE O2 Supply

- 1-Increase diastolic time: B-blocker
- 2-Decrease coronary tone: nitrate, ca-blocker *Dilation*
- 3-Decrease LV diastolic pressure: nitrate
- 4-Correct coronary stenosis: PCI, CABG
- 5-Increase O2 capacity of blood: transfusion if anemia

DECREASE O2 Demand

- 1-Decrease heart rate: B-blocker, ca-blocker
- 2-Decrease contractility: B-blocker, ca-blocker
- 3- Decrease wall tension (LV pressure and cavity radius): nitrate
- 4- metabolic: trimetazidine

β -blocker
 \uparrow diastole time
 \downarrow HR
 \downarrow contractility

Ca-blocker
 \downarrow coronary tone
 \downarrow HR
 \downarrow contractility

nitrate
 \downarrow coronary tone
 \downarrow LV diastolic pressure
 \downarrow LV wall tension

Treatment in practice

1-General measures

2-Aspirin $\xrightarrow{\text{sublingual}}$ according to chest pain frequency

3-Nitrate: S/L, Oral, dermal

3-B-blocker

4-Statins: LDL > 100 mg/dl (70mg/dl)

5-Ca-blocker

6-Angio :PTCA,CABG

New medical and invasive therapies for refractory angina

Inhibition of fatty acid metabolism: **trimetazidine**

Potassium channel activators: **Nicorandil**.

Ranolazine: interact with sodium channel

Ivabradine: SA inhibitor

Endothelin Receptor Blockers: **bosentan**

Testosterone: improve endoth dysfunction.

Enhanced external balloon counterpulsation

Spinal cord stimulation.

Laser revascularization, angiogenesis.

Prognostic Indicators of Coronary Artery Disease:

1- Left Ventricular Function:

Normal: 50-75%

< 50% associated with increased mortality

2- Vessel(s) involved: severity and extent of ischemia

mortality/year

2% single vessel-----12% left main stem

Differential diagnosis of angina

- 1- Neuromuscular disorder
- 2- Respiratory disorders
- 3-Upper GI disorder
- 4- Psychological
- 5- Syndrome X

Cardiac Syndrome X

↓ vasodilation on exercise

Typical, exertional angina with positive exercise stress test

normal
coronary

Anatomically normal coronary arteries

Reduced capacity of vasodilation in microvasculature

F>M

Young > Elderly

Excellent prognosis

Antianginal therapy is rarely effective

Long term prognosis very good

Case History

A 45 Year old male, presented with recurrent attacks of chest pain last few minutes, during attacks of chest pain the ECG abnormalities as attached ECG. After pain subsides the ECG back to normal.

What is the diagnosis?


Intermediate -----

ST elevation


V1, V5, V6

after nitrate → relieved

ECG



After sublingual isosorbid dinitrate tablet



VARIANT ANGINA-PRINZMETAL ANGINA

Chest pain with ST-Segment elevation

Usually at rest, at night

Troponin: negative

Female > male

Spasm of large epicardial coronary vessel during the attack
transmural ischemia

70% on top of atherosclerosis

Vasospastic symptoms in other organs: Migraine, rhynauds

Can cause arrhythmias and death

Treatment: CA-blocker, Nitrate → β B blocks β -receptors & leave α -receptors
+ cause vasoconstriction

B-blocker is contraindicated

Prognosis: 5 year mortality < 5%