

Pericardium

Hanna Al-Makhamreh, MD.FACC
Associate Professor of Cardiology
University of Jordan


Edited by: Ruaa Adeib

Pericardium


- **Introduction:**
- The pericardium is a fibroelastic sac made up of visceral and parietal layers separated by a space, the pericardial cavity.
- In healthy individuals, the pericardial cavity contains **15-50 mL** of an ultrafiltrate of plasma.

↳ heart is fixed in mediastinum

Pericardium



Pericardium



Function

- The pericardium acts as **mechanical protection for the heart and big vessels, anchoring the heart in the mediastinum and lubrication to reduce friction between the heart and the surrounding structures**

Pericarditis

- Acute pericarditis refers to inflammation of the pericardial sac.

pericardium alone or with myocarditis ← → predominantly affect the myocardium with pericardial involvement

- The term myopericarditis, or perimyocarditis, is used for cases of acute pericarditis that also demonstrate myocardial inflammation.

Pericarditis

viral / idiopathic

- Most cases of acute pericarditis are considered of possible or confirmed viral origin, although the exact etiology of most cases remains undetermined following a traditional diagnostic approach.
- Acute pericarditis is a common disorder in several clinical settings, where it may be the first manifestation of an underlying systemic disease or may represent an isolated process.
Malignancy can present as pericarditis but mostly as pericardial effusions

Major causes of pericardial disease

Idiopathic

In most case series, the majority of patients are not found to have an identifiable cause of pericardial disease. Frequently such cases are presumed to have a viral or autoimmune etiology.

Infections

- Viral - Coxsackievirus, echovirus, adenovirus, EBV, CMV, influenza, varicella, rubella, HIV, hepatitis B, mumps, parvovirus B19, vaccinia (smallpox vaccination)
- Bacterial - Staphylococcus, Streptococcus, pneumococcus, Haemophilus, Neisseria (gonorrhoeae or meningitidis), Chlamydia (pneumonia or trachomatis), Legionella, tuberculosis, Salmonella, Lyme disease
- Mycoplasma
- Fungal - Histoplasmosis, aspergillosis, blastomycosis, coccidioidomycosis, actinomycosis, nocardia, candida
- Parasitic - Echinococcus, amebiasis, toxoplasmosis
- Infective endocarditis with valve ring abscess

Radiation

Neoplasm

higher prevalence in cancers

اکثر شایع نوع کالج

- Metastatic - Lung or breast cancer, Hodgkin's disease, leukemia, melanoma
- Primary - Rhabdomyosarcoma, teratoma, fibroma, lipoma, leiomyoma, angioma
- Paraneoplastic

Cardiac

- Early infarction pericarditis
- Late postcardiac injury syndrome (Dressler's syndrome), also seen in other settings (eg. post-myocardial infarction and post-cardiac surgery)
- Myocarditis
- Dissecting aortic aneurysm

Trauma

- Blunt
- Penetrating
- Iatrogenic - Catheter and pacemaker perforations, cardiopulmonary resuscitation, post-thoracic surgery

Autoimmune

- Rheumatic diseases - Including lupus, rheumatoid arthritis, vasculitis, scleroderma, mixed connective disease
- Other - Wegener's granulomatosis, polyarteritis nodosa, sarcoidosis, inflammatory bowel disease (Crohn's, ulcerative colitis), Whipple's, giant cell arteritis, Behcet's disease, rheumatic fever

Drugs

- Procainamide, isoniazid, or hydralazine as part of drug-induced lupus
- Other - Cromolyn sodium, dantrolene, methysergide, anticoagulants, thrombolytics, phenytoin, penicillin, phenylbutazone, doxorubicin

Metabolic

- Hypothyroidism - Primarily pericardial effusion
- Uremia
- Ovarian hyperstimulation syndrome

Adapted from: Shabetai R. Diseases of the pericardium. In: Hurst's The Heart, 8th ed, Schifert RC, Alexander RW, et al (Eds).

Pericarditis

upper/flu like illness

- **Clinical presentation:**
- Acute pericarditis can present in a variety of ways, depending on the underlying etiology.
- Patients with an infectious etiology may present with signs and symptoms of systemic infection such as fever and leukocytosis.
- Viral etiologies in particular may be preceded by “flu-like” respiratory or gastrointestinal symptoms.

Pericarditis

- Patients with a known autoimmune disorder or malignancy may present with signs or symptoms specific to their underlying disorder.
- The major clinical manifestations of acute pericarditis include:

Pericarditis

- **Chest pain** — typically sharp and pleuritic, improved by sitting up and leaning forward. *↑ with supine*
- **Pericardial friction rub** — a superficial scratchy or squeaking sound best heard with the diaphragm of the stethoscope over the left sternal border.
- **Electrocardiogram (ECG) changes** — new widespread ST elevation or PR depression
- **Pericardial effusion.** *Echo*

Pericarditis

- **I-Chest pain:** بعد نكته HI نكته sudden
- The vast majority of patients with acute pericarditis present with chest pain (>95% of cases). جمع ايجاد trabisious
- Chest pain that results from acute pericarditis is typically fairly **sudden** in onset and occurs over the **anterior chest**.
- Unlike pain from myocardial ischemia, chest pain due to pericarditis is most often sharp and pleuritic in nature, **with exacerbation by inspiration or coughing.**

Pericarditis

- One of the most distinct features is the **tendency for a decrease in intensity when the patient sits up and leans forward.**
- This position (seated, leaning forward) tends to reduce pressure on the parietal pericardium, particularly with inspiration, and may also allow for splinting of the diaphragm.

Pericarditis

- **II-Pericardial friction rub:**
- The presence of a pericardial friction rub on physical examination is highly specific for acute pericarditis.
- Pericardial friction rubs, which occur during the maximal movement of the heart within its pericardial sac, are said to be generated by **friction between the two inflamed layers of the pericardium.**

Pericarditis

- Pericardial rubs have a superficial scratchy or squeaking quality that is best heard with the diaphragm of the stethoscope.
- They may be localized or widespread, but are usually loudest over the left sternal border. *at end of exhalation*

ECG

- **III-Electrocardiogram:**
- Changes in the electrocardiogram (ECG) in patients with acute pericarditis signify inflammation of the epicardium, since the parietal pericardium itself is electrically inert.

ECG

MI → convex downward
concave upward


- The typical progression of ECG changes in patients with acute pericarditis is described below:
- **Stage 1**, seen in the **first hours to days**, is characterized by **diffuse ST elevation** (typically concave up) with **reciprocal ST depression in leads aVR and V1**.
- There is also an atrial current of injury, reflected by **elevation of the PR segment in lead aVR** and **depression of the PR segment in other leads**.

ECG

- **Stage 2**, typically seen in the first week, is characterized by normalization of the ST and PR segments.
- **Stage 3**, is characterized by the development of diffuse T wave inversions, generally **after** the ST segments have become isoelectric.
- **Stage 4**, is represented by normalization of the ECG or indefinite persistence of T wave inversions ("chronic" pericarditis).

Most common cause of effusion is pericarditis


Normal ECG



Normal sinus rhythm at a rate of 71 beats/min, a P wave axis of 45° , and a PR interval of 0.15 sec.

Courtesy of Morton Arnsdorf, MD.

Pericarditis



Electrocardiogram in acute pericarditis showing diffuse upsloping ST segment elevations seen best here in leads II, III, aVF, and V2 to V6. There is also subtle PR segment deviation (positive in aVR, negative in most other leads). ST segment elevation is due to a ventricular current of injury associated with epicardial inflammation; similarly, the PR segment changes are due to an atrial current of injury which, in pericarditis, typically displaces the PR segment upward in lead aVR and downward in most other leads.

Courtesy of Ary Goldberger, MD.

Characteristic	Pericardial Pain	Myocardial Ischemic Pain
Quality	Sharp, pleuritic	Pressure, heaviness, tightness, constricting
Location	Left precordial or retrosternal	Retrosternal
Radiation pattern	Left trapezius ridge	Left shoulder, left arm
Duration	Hours or days	1-15 minutes (angina) >20 minutes (unstable angina) Hours (myocardial infarction)
Relation to exercise	Unrelated	Related (stable angina) Unrelated (unstable angina or myocardial infarction)
Relation to position	Relieved by leaning forward Aggravated by assuming a recumbent position	Unrelated
ECG findings	Initial ECG changes that accompany onset of chest pain: ST-segment elevation that is upwardly concave and diffuse (occurs in all leads except aVR and V ₁) T waves invert after ST-segment elevation resolves and not associated with loss of R-wave voltage or Q waves (occurs several days after onset of chest pain) PR-segment depression present in 80% (occurs in all limb and precordial leads except aVR); reciprocal PR-segment elevation may occur in aVR. Q waves absent	ST-segment elevation is downwardly concave and localized; or ST-segment depression T waves invert while ST-segment elevation is present. May be associated with loss of R-wave voltage or appearance of Q waves PR-segment depression rarely present Q waves may be present
Echocardiographic findings	No left ventricular regional wall motion abnormality	Left ventricular regional wall motion abnormality in distribution of coronary artery

Diagnosis

- Echocardiography is often normal in patients with the clinical syndrome of acute pericarditis unless there is an associated pericardial effusion.
- Chest radiography is typically normal in patients with acute pericarditis. Although patients with a substantial pericardial effusion may exhibit an enlarged cardiac silhouette with clear lung fields .

global hypokinesia

Diagnosis

- Acute pericarditis may be associated with increases in serum biomarkers of myocardial injury such as cardiac troponin or MB fraction of creatine kinase.
- Since pericarditis is an inflammatory disease, laboratory signs of inflammation are common in patients with acute pericarditis.
- These include elevations in the white blood cell count, **ESR** erythrocyte sedimentation rate, and serum C-reactive protein concentration.

Diagnostic criteria for acute pericarditis and myopericarditis in the clinical setting

Acute pericarditis (at least 2 criteria of 4 should be present)*:

1. Typical chest pain
2. Pericardial friction rub
3. Suggestive ECG changes (typically widespread ST segment elevation)
4. New or worsening pericardial effusion

Myopericarditis:

1. Definite diagnosis of acute pericarditis, **PLUS**
2. Suggestive symptoms (dyspnea, palpitations, or chest pain) and ECG abnormalities beyond normal variants, not documented previously (ST/T abnormalities, supraventricular or ventricular tachycardia or frequent ectopy, atrioventricular block), **OR** focal or diffuse depressed LV function of uncertain age by an imaging study.
3. Absence of evidence of any other cause
4. One of the following features: evidence of elevated cardiac enzymes (creatine kinase-MB fraction, or troponin I or T), **OR** new onset of focal or diffuse depressed LV function by an imaging study, **OR** abnormal imaging consistent with myocarditis (MRI with gadolinium, gallium-67 scanning, anti-myosin antibody scanning)

Case definitions for myopericarditis include:

Suspected myopericarditis: criteria 1 plus 2 and 3

Probable myopericarditis: criteria 1, 2, 3, and 4

Confirmed myopericarditis[†]: histopathologic evidence of myocarditis by endomyocardial biopsy or on autopsy

* Pericardial effusion confirms the clinical diagnosis but its absence does not exclude it.

† In clinical practice a confirmed diagnosis would require an endomyocardial biopsy that is not warranted in self-limited cases with predominant pericarditis.

Reproduced with permission from: Imazio M, Trinchero R. Triage and management of acute pericarditis. *Int J Cardiol* 2006, doi:10.1016/j.ijcard.2006.07.100. Copyright: © 2006 Elsevier.

Treatment

- In cases of pericarditis due to an identifiable cause (eg, bacterial infection or malignancy, autoimmune), management is focused upon the underlying disorder and, if necessary, drainage of an associated pericardial effusion percutaneously or surgically .

Treatment

- For most patients with acute idiopathic or viral ^{3 months} pericarditis, combination therapy with colchicine plus NSAIDs rather than NSAIDs ^{2-4 weeks} alone.
- This is based upon a "reduced rate of recurrent pericarditis" and a low incidence of side effects with colchicine. ↓ severity

Treatment

- In patients with acute pericarditis following an MI, aspirin plus colchicine rather than another NSAID plus colchicine
- This is principally due to the possibility that other NSAIDs may interfere with healing and scar formation.
- Although the evidence of potential harm from glucocorticoids and NSAIDs other than aspirin is modest, there is no evidence that these medications improve outcomes.

rather than any
other NSAID

post MI → aspirin

Treatment

- *For these reasons glucocorticoids and NSAIDs other than aspirin should generally be AVOIDED in patients with acute pericarditis following an acute MI.*

Lupus / Rheumatoid must take

uremia → hemodialysis

Steroids

↓
→ second line therapy (if NSAIDs didn't work)

- pericarditis due to connective tissue disease, autoreactive (immune-mediated) pericarditis, uremic pericarditis not responding to dialysis, and to patients who have contraindications to NSAID therapy.
- Glucocorticoid therapy is also used for patients with idiopathic or viral pericarditis that is refractory to combination therapy with NSAIDs and colchicine.

Recurrent Pericarditis

- Recurrent pericarditis may occur in up to 30% of patients after an initial episode of acute pericarditis. Treatment should consist of an NSAID, typically with a 2- to 4-week taper after the resolution of symptoms, along with at least 6 months of colchicine (with weight-adjusted dosing).
- Steroids
- Anti-interleukin 1 therapy (e.g., anakinra and rilonacept)
- Azathioprine, methotrexate, mycophenolate mofetil, and intravenous immunoglobulins.

Pericarditis

Drug therapy in acute pericarditis

Drug	Dose	Duration of therapy	Tapering
For initial combination treatment of most patients:			
NSAID Ibuprofen	400 to 800 mg three times daily	1 to 2 weeks	Decrease the dose weekly
OR			
NSAID Indomethacin	50 mg three times daily	1 to 2 weeks	Decrease the dose weekly
PLUS			
Colchicine	0.5 mg two times daily	3 months	Usually not tapered
For initial combination therapy of patients following myocardial infarction:			
Aspirin	650 to 1000 mg three times daily	1 to 2 weeks	Decrease the dose weekly
PLUS			
Colchicine	0.5 mg two times daily	3 months	Usually not tapered
For refractory cases or patients with a contraindication to NSAID therapy:			
Prednisone	0.2 to 0.5 mg/kg/day	2 weeks	Slow tapering, see text
PLUS			
Colchicine	0.5 mg two times daily	3 months	Usually not performed

High Risk Features

1-High fever ($>38^{\circ}\text{C}$)

2-Subacute onset

3-Large pericardial effusion or tamponade


4-Lack of response to anti-inflammatory therapy after 1 week of treatment

5-Evidence for myocardial involvement

Additional predictors of elevated risk include immunosuppression, oral anticoagulation, and trauma.

إذا ما عنده هذه الأشياء، بترجمه

Initial clinical and echocardiographic evaluation of patients with suspected acute pericarditis




Reproduced with permission and modified from: Imazio, M, Demichelis, B, Parrini, I, et al. Day-hospital treatment of acute pericarditis: a management program for outpatient therapy. *J Am Coll Cardiol* 2004; 43:1042. Copyright © 2004 Elsevier.

Pericardial Effusion/Tamponade

- Causes
 - Trauma, uremia, anticoagulation, neoplasm, infection
- Clinical signs (Beck's triad, 1935)
 - Hypotension, JVD, muffled heart sounds
 - pulsus paradoxus *drop in systolic BP of 10 mm Hg during inspiration*
acute bronchitis
- ECG
 - Electrical alternans (beat to beat alteration in the amplitude of the QRS complex)
 - low voltage
- ECHO findings
 - Effusion
 - RV diastolic collapse (specific for tamponade)

Pericardial Effusion



electrical alternans

From ECGs for the Emergency Physician Vol 1 (Matta, Brady, Blackwell, 2003)

Tamponade

- Dyspnea
- Tachycardia
- JVD
- Hypotension
- Pulsus paradoxus >10 mm Hg supports dx

Echo Findings:

- Effusion
- Diastolic collapse of right-sided chambers
- Increased respiratory variation of peak inflow velocities through TV and MV
- Dilated IVC without respire-phasic variation

Management

Hemodynamically stable:

IVF, close monitoring, serial TTE, treat underlying cause

Unstable:

-Aggressive IVF


-Pericardiocentesis or surgery

intra-aortic balloon pump

-IABP for refractory hypotension

-Minimize PEEP

Electrical alternans



Sinus tachycardia with electrical alternans which is characterized by beat-to-beat alternation in the QRS appearance (best seen in leads V2 to V4). These findings are strongly suggestive of pericardial effusion, usually with cardiac tamponade. The alternating ECG pattern is related to back-and-forth swinging motion of the heart in the pericardial fluid. *Courtesy of Ary Goldberger, MD.*

Chest x-ray of a pericardial effusion



Cardiomegaly due to a massive pericardial effusion. At least 200 mL of pericardial fluid must accumulate before the cardiac silhouette enlarges.

Courtesy of Massimo Imazio, MD, FESC.

Subcostal view

Sternum

Effusion

Effusion

RA


RV

LV

LA

Pericardial
effusion

Thoracic wall



Hub of needle
angled approximately
15 degrees above skin

Apical
area


Paraxiphoid
area

Constrictive Pericarditis

- Thick dense fibrous obliteration with calcification of the pericardial sac encasing the heart
- limiting diastolic expansion and restricting cardiac output

uncommon disease / very challenging

Constrictive Pericarditis



Absence of calcification does not r/o constrictive pericarditis

x-ray

Constrictive Pericarditis

- Evidence of right heart failure
- Kussmaul sign: no fall or even elevation JVP with inspiration
- Abnormal echo

Constrictive Pericarditis

- Dyspnea
- Fatigue
- JVD
- Hepatomegaly and ascites
- Edema
- Neck veins distend with inspiration (Kussmaul's sign)
- Pericardial knock (early diastolic sound)
- Afib in 20%
- Cold leg, no palpable pulse

Causes:

- Cardiac surgery
- Viral infection
- Acute pericarditis
- Mediastinal irradiation
- TB
- Rheumatoid arthritis, CTD

Management

- * Diurese with caution
- * Rate-control with caution
- * 2-3 month trial of conservative measures prior to pericardiectomy

Pericardiectomy indicated in:

NYHA Class II or III with persistent symptoms

In NYHA Class IV and those with cachexia, cirrhosis, markedly reduced cardiac output, benefit of surgery is negligible


Table 10 Constrictive pericarditis vs. restrictive cardiomyopathy: a brief overview of features for the differential diagnosis (Modified from Imazio et al.⁵¹)

Diagnostic evaluation	Constrictive pericarditis	Restrictive cardiomyopathy
Physical findings	Kussmaul sign, pericardial knock	Regurgitant murmur; Kussmaul sign may be present, S3 (absent).
ECG	Low voltages, non-specific ST/T changes, atrial fibrillation	Low voltages, pseudoinfarction, possible widening of QRS, left-axis deviation, atrial fibrillation
Chest X-ray	Pericardial calcifications (1/3 of cases)	No pericardial calcifications
Echocardiography	<ul style="list-style-type: none"> • Segal bounce. • Pericardial thickening and calcifications. • Respiratory variation of the mitral peak E velocity of >25% and variation in the pulmonary venous peak D flow velocity of >20%. • Colour M-mode flow propagation velocity (Vp) >45 cm/sec. • Tissue Doppler: peak e' >8.0 cm/s. 	<ul style="list-style-type: none"> • Small left ventricle with large atria, possible increased wall thickness. • E/A ratio >2, short DT. • Significant respiratory variations of mitral inflow are absent. • Colour M-mode flow propagation velocity (Vp) <45 cm/sec. • Tissue Doppler: peak e' <8.0 cm/s.
Cardiac Catheterization	Dip and plateau/ or 'square root' sign, right ventricular diastolic and left ventricular diastolic pressures usually equal, ventricular interdependence (i.e. assessed by the systolic area index >1.1) ⁴	Marked right ventricular systolic hypertension (>50 mmHg) and left ventricular diastolic pressure exceeds right ventricular diastolic pressure (LVEDP >RVEDP) at rest or during exercise by 5 mmHg or more (RVEDP <1/3 RVSP).
CT/MR	Pericardial thickness >3-4 mm, pericardial calcifications (CT), ventricular interdependence (real-time cine CMR).	Normal pericardial thickness (<3.0 mm), myocardial involvement by morphology and functional study (CMR).

Case 1


- A 46-year-old-male patient with a pmhs of hypertension and smoking presented to the ED with chest pain.


Name: 12-Lead 2 HR 66 bpm • ***** ACUTE MI SUSPECTED ***** • Lateral injury pattern
 IL: [REDACTED] 02:49:43 • **Abnormal ECG **Unconfirmed****
 Patient ID: PR 0.156s QRS 0.208s • Normal sinus rhythm
 Incident: QT/QTc 0.336s/0.415s • Left atrial enlargement
 Age: 71 Sex: P-QRS-T Axes 72° 5° 12° • Anteroseptal infarct, possibly acute



Case 2

- A 25-year old male pt. presented with worsening dyspnea to the ED. His heart rate was 140 bpm and BP 80/40 mm Hg.





Thank you