



Blood and Blood Product Transfusion

Amjad Bani Hani

Assistant Prof. of Cardiac Surgery and Intensive Care

The University Of Jordan

- Blood and Blood Products
- Indications for transfusion
- Complications of transfusion
- Massive blood transfusion

- **Approximately 15 million red blood cell (RBC) units are transfused annually in the United States (1)**

1. U.S. Department of Health and Human Services. The 2009 national blood collection and utilization survey report. Washington, DC: U.S. Department of Health and Human Services, Office of the Assistant Secretary for Health; 2011.

- About 85 million are transfused annually worldwide.


Takei T, Amin NA, Schmid G, Dhingra-Kumar N, Rugg D. Progress in global blood safety for HIV. *J Acquir Immune Defic Syndr.* 2009;52 Suppl 2:S127-31. [PMID: 19901625]

- Approximately 118 million units of blood are collected worldwide each year.¹

World Health Organization. Blood transfusion. June 8, 2022. Accessed February 23, 2023. <https://www.who.int/news-room/facts-in-pictures/detail/blood-transfusion>

- Clinicians should offer RBC transfusion to patients only when benefits outweigh harms

Figure. Adverse effects of RBC transfusion contrasted with other risks.



RATIONALE FOR TRANSFUSION

- **Role of blood in oxygen delivery**
- **Impact of anemia on morbidity and mortality**


Why?

- The body at rest uses approx **250ml O₂/L blood**
- O₂ delivery can fall with a reduction in any of:
 - Cardiac Output
 - Hb concentration
 - O₂ saturation
- Organs most sensitive to hypoxia are Heart and Brain



Why?

- The purpose of a red cell transfusion is to improve the oxygen carrying capacity of the blood.
- Oxygen delivery to tissues (O₂ Flux)
= Cardiac Output x Oxygen content of blood




Blood Transfusion in the Operating Room Is Bad!

- Cardiac Surgery
- Thoracic operations
- Vascular operations
- Cancer procedures
- General Surgery
- Cardiology doesn't get a pass!
 - PCI outcomes worse w/ blood transfusion




Transfusion & Serious Morbidity in 4,445 Cardiac Surgical Patients



- Serious morbidity and mortality increase with the amount transfused.



Intraoperative Blood Transfusion & Lung Surgery




Ferraris, 2011



Blood Transfusion in General Surgical Population

Intraoperative Blood Transfusion and NSQIP Surgical Outcomes in 941,496 Patients






Blood Components

- Prepared from Whole blood collection or apheresis
- Whole blood is separated by differential centrifugation
 - Red Blood Cells (RBC's)
 - Platelets
 - Plasma
 - Cryoprecipitate
 - Others
- Others include Plasma proteins—IVIg, Coagulation Factors, albumin, Anti-D, Growth Factors, Colloid volume expanders
- Apheresis may also used to collect blood components




Differential Centrifugation

First Centrifugation



Differential Centrifugation

Second Centrifugation



Whole Blood



- Storage
 - 4° for up to 35 days
- Indications
 - Massive Blood Loss/Trauma
- Considerations
 - Use filter as platelets and coagulation factors will not be active after 3-5 days
 - Donor and recipient must be ABO identical



RBC Concentrate

- Storage
 - 4° for up to 42 days, can be frozen
- Indications
 - Many indications—ie anemia, hypoxia, etc.
- Considerations
 - Recipient must not have antibodies to donor RBC's (note: patients can develop antibodies over time)
 - Usual dose 10 cc/kg (will increase Hgb by 2.5 gm/dl)
 - Usually transfuse over 2-4 hours (slower for chronic anemia)

Platelets



- Storage
 - Up to 5 days at 20-24°
- Considerations
 - Contain Leukocytes and cytokines
 - 1 unit/10 kg of body weight increases Plt count by 50,000
 - Donor and Recipient must be ABO identical

- Indications
- 10,000/mm³ in stable, non-bleeding patients,
- 20,000/mm³ in unstable non-bleeding patients
- 50,000/mm³ in patients undergoing invasive procedures or actively bleeding.

- Neurologic or ophthalmologic or Cardiac procedures require a platelet count near 100,000/mm³.

FFP

- Contents—Coagulation Factors (1 unit/ml)
- Storage
 - FFP--12 months at –18 degrees or colder
- Indications
 - Coagulation Factor deficiency, fibrinogen replacement, DIC, liver disease, exchange transfusion, massive transfusion

- Considerations

- Plasma should be recipient RBC ABO compatible

- Usual dose is 20 cc/kg to raise coagulation factors approx 20%

Cryoprecipitate

- Description
 - Precipitate formed/collected when FFP is thawed at 4°
- Storage
 - After collection, refrozen and stored up to 1 year at -18°
- Indication
 - Fibrinogen deficiency or dysfibrinogenemia
 - vonWillebrands Disease
 - Factor VIII or XIII deficiency
 - DIC

- Considerations
 - ABO compatible preferred (but not limiting)
 - Usual dose is 1 unit/5-10 kg of recipient body weight

Leukocyte Reduction Filters

- Used for prevention of transfusion reactions
- Filter used with RBC's, Platelets, FFP, Cryoprecipitate
- May reduce RBC's by 5-10%
- Does not prevent Graft Verses Host Disease (GVHD)

When to transfuse

Background

- Carson et al. “Mortality and morbidity in patients with very low **postoperative** Hb levels who decline blood transfusion.”
Transfusion 2002
 - Mortality
 - Hgb 7.1 to 8.0 (n = 99) — zero percent
 - Hgb 5.1 to 7.0 (n = 110) — 9 percent
 - Hgb 3.1 to 5.0 (n = 60) — 30 percent
 - Hgb ≤ 3.0 (n = 31) — 64 percent

The TRICC Study

- Enrolled 838 euvolemic, anemic, critically ill pts who were admitted to 1 of 25 Canadian ICUs
- Patients were stratified according to center and disease severity (APACHE II) and placed into one of two groups
 - ❑ **Restrictive group**: Transfuse if Hb < 7 and maintain between 7 and 9
 - ❑ **Liberal group**: Transfuse if Hb < 10 and maintain between 10 and 12
- The primary outcome measure was death from all causes in the 30 days after randomization


The TRICC Study

No difference 30 day mortality


In “healthy” (APACHE II < 20) and young (<55yrs) patients

Transfusion increased mortality

The TRICC Study



8.7% vs 16.1%



5.7% vs 13.0%


“A restrictive red blood cell transfusion strategy generally appears to be safe in most critically ill patients with cardiovascular disease... with the possible exception of patients with acute myocardial infarction and unstable angina.”

CRIT Study


- Prospective, multiple center, observational cohort study of **4,892 ICU pts** in the US
- Propensity score matched
- Designed to examine the relationship of anemia and RBC transfusion with clinical outcomes
- Almost 95% of patients admitted to the ICU have a Hb level below “normal” by day 3
- In total, 11,391 RBC units were transfused.
- Overall, 44% of pts admitted to the ICU received one or more RBC units while in the ICU

CRIT Results

35% of Blood transfused in patients with Hgb ≥ 9



The mean pre-transfusion Hb was 8.6 ± 1.7 g/dL



RBC transfusion was independently associated with higher mortality (OR 1.65 CI 1.35-2.03). OR 2.62 if 3-4 units transfused p < 0.0001

When to Pull the “transfusion Trigger?”

- **Should not be based solely on hemoglobin number.**
- Decision should consider clinical scenario, patient characteristics, and symptoms.

When to Pull the “transfusion Trigger?”

American Association of Blood Banks Guidelines

- Hgb <7
- Hgb 7-8
- Transfusion recommended
- **Restrictive Transfusion Strategy** for stable patients
Consider transfusion only if post-operative or symptomatic (chest pain, orthostatic hypotension or tachycardia unresponsive to fluid resuscitation, or congestive heart failure).

– Hgb 8 – 10

- TRANSFUSION GENERALLY NOT INDICATED

Can consider Tx in special circumstances (ie ACS w/ active ischemia, symptomatic anemia, active bleeding).

– Hgb >10

- TRANSFUSION NOT INDICATED

Transfusion Complications

- Acute Transfusion Reactions (ATR's)
- Chronic Transfusion Reactions
- Transfusion related infections



Acute Transfusion Reactions

- Hemolytic Reactions (AHTR)
- Febrile Reactions (FNHTR)
- Allergic Reactions
- TRALI
- Coagulopathy with Massive transfusions
- Bacteremia

Acute Hemolytic Transfusion Reactions (AHTR)

- Occurs when incompatible RBC's are transfused into a recipient who has pre-formed antibodies (usually ABO or Rh)
- Antibodies activate the complement system, causing intravascular hemolysis
- Symptoms occur within minutes of starting the transfusion
- This hemolytic reaction can occur with as little as 1-2 cc of RBC's
- Labeling error is most common problem
- **Can be fatal**

Symptoms of AHTR

- High fever/chills
- Hypotension
- Back/abdominal pain
- Oliguria
- Dyspnea
- Dark urine
- Pallor

What to do?

If an AHTR occurs



- **STOP TRANSFUSION**
- **ABC's**
- Maintain IV access and run IVF (NS or LR)
- Monitor and maintain BP/pulse
- Give diuretic
- Obtain blood and urine for transfusion reaction workup
- Send remaining blood back to Blood Bank

Blood Bank Work-up of AHTR

- Check paperwork to assure no errors
- Check plasma for hemoglobin
- DAT (Direct Antiglobulin Test)
- Repeat crossmatch
- Repeat Blood group typing
- Blood culture



Labs found with AHTR

- Hemoglobinemia
- Hemoglobinuria
- Positive DAT
- Hyperbilirubinemia
- Abnormal DIC workup

Monitoring in AHTR

- Monitor patient clinical status and vital signs
- Monitor renal status (BUN, creatinine)
- Monitor coagulation status (DIC panel– PT/PTT, fibrinogen, D-dimer/FDP, Plt, Antithrombin-III)
- Monitor for signs of hemolysis (LDH, bili, haptoglobin)

Febrile Nonhemolytic Transfusion Reactions (FNHTR)

- Definition--Rise in patient temperature $>1^{\circ}\text{C}$ (associated with transfusion without other fever precipitating factors)
- Occurs with approx 1% of PRBC transfusions and approx 20% of Plt transfusions
- FNHTR caused by alloantibodies directed against HLA antigens

What to do?

If an FNHTR occurs

- STOP TRANSFUSION
- Use of Antipyretics
- Use of Corticosteroids for severe reactions
- Use of Narcotics for shaking chills
- Future considerations
 - May prevent reaction with leukocyte filter
 - Use single donor platelets
 - Washed RBC's or platelets

Allergic Nonhemolytic Transfusion Reactions


- Etiology
 - May be due to plasma proteins or blood preservative/anticoagulant
 - Best characterized with IgA given to an IgA deficient patients with anti-IgA antibodies
- Presents with urticaria and wheezing
- Treatment
 - Mild reactions—Can be continued
 - Severe reactions—Must STOP transfusion and may require steroids or epinephrine
- Prevention—Premedication (Antihistamines)

TRALI

Transfusion Related Acute Lung Injury

- Clinical syndrome similar to ARDS
- Occurs 1-6 hours after receiving plasma-containing blood products
- Caused by WBC antibodies present in donor blood that result in pulmonary leukostasis
- Treatment is supportive
- High mortality

Initial approach to a suspected acute transfusion reaction



This graphic includes some of the most common and life-threatening reactions; other reactions are also possible and should be pursued if the clinical picture seems inconsistent with one of these. The transfusion service should be notified of any severe transfusion reaction and may request samples of the transfused product and patient blood; the transfused product should not be discarded until discussion with the transfusion service has taken place. In cases of suspected AHTR, the transfusion service must be contacted immediately because another patient may be at risk of receiving the incorrect blood product. Refer to UpToDate topics on transfusion reactions for further details of the evaluation and management of these conditions.

TACO: transfusion-associated circulatory overload; TRALI: transfusion-related acute lung injury; FNHTR: febrile nonhemolytic transfusion reaction; AHTR: acute hemolytic transfusion reaction; ALI/ARDS: acute lung injury/adult respiratory distress syndrome; DAT: direct antiglobulin test (Coombs test); CBC: complete blood count; CXR: chest x-ray; LDH: lactate dehydrogenase; DIC: disseminated intravascular coagulation.

Massive Transfusions

- Coagulopathy may occur after transfusion of massive amounts of blood (trauma/surgery)
- Coagulopathy is caused by failure to replace plasma
- Electrolyte abnormalities
 - Due to citrate binding of Calcium
 - Also due to breakdown of stored RBC's

Bacterial Contamination

- More common and more severe with platelet transfusion (platelets are stored at room temperature)
- Organisms
 - Platelets—Gram (+) organisms, ie Staph/Strep
 - RBC's—Yersinia, enterobacter
- Risk increases as blood products age (use fresh products for immunocompromised)

Chronic Transfusion Reactions

- Alloimmunization
- Transfusion Associated Graft Verses Host Disease (GVHD)
- Iron Overload
- Transfusion Transmitted Infection



Transfusion Associated Infections

- Hepatitis C
- Hepatitis B
- HIV
- CMV
 - CMV can be diminished by leukoreduction, which is indicated for immunocompromised patients



Risks of Transfusion: Infectious Disease

- ✓ HIV = 1 in 1.8 million
- ✓ HCV = 1 in 1.6 million
- ✓ HBV = 1 in 220,000

HIV = human immunodeficiency virus.

HCV = hepatitis C virus.

HBV = hepatitis B virus.


Busch MP, et al. *JAMA*. 2003;289:959-62.

**Why is blood transfusion
NOT associated with
improved outcome?**


Stored RBCs

- Decreased RBC deformability
- Decreased 2,3, DPG
- Metabolic acidosis
- Altered oxygen carrying capacity
- Increased red cell death with increased age of blood (~30% dead)
- No improvement in oxygen utilization at the tissue level


Age of Blood



Day 1



Day 21






Day 35

Scanning electron micrographs of red blood cells isolated from stored blood on Day 1, Day 21, and Day 35. During storage, the shape of RBCs changed gradually from normal discoid to echinocytes (dented or shriveled red cells).


Reproduced with permission from: Hovav et al. *Transfusion*. 1999;39:277-281.

Poor Efficacy of Blood Tx

- RBCs stored > 15 days lose deformability and ATP
- Altered capillary lumen size (decreased cross-sectional diameter) in critically ill patients
- Increased “stickiness” (adherence) of RBCs to altered endothelium in the microcirculation of critically ill pts.



Distribution of Transfused Units by Age of Blood - CRIT Study



60% of Blood transfused is
> 20 days old




Massive blood Transfusion



Causes of death following multiple trauma

Etiology of trauma deaths

■ CNS ■ Hemorrhage ■ MOF



Classification of Hemorrhage

American College of Surgeons Committee on Trauma
Advanced Trauma Life Support Program

	CLASS I	CLASS II	CLASS III	CLASS IV
Blood loss (ml)	Up to 750	750-1,500	1,500-2,000	≥ 2,000
Blood loss (% blood volume)	Up to 15%	15%-30%	30%-40%	≥40%
Pulse rate	<100	>100	>120	≥140
Blood pressure	Normal	Normal	Decreased	Decreased
Pulse pressure (mmHg)	Normal or increased	Decreased	Decreased	Decreased
Capillary refill test	Normal	Positive	Positive	Positive
Respiratory rate	14-20	20-30	30-40	>35
Urine output (ml/hr)	≥30	20-30	5-15	Negligible
CNS — mental status	Slightly anxious	Mildly anxious	Anxious, confused	Confused, lethargic

What is Massive transfusion?

10 units of red cells in 24 hours

Total blood volume is replaced within 24 hours

Three units over one hour with ongoing bleeding

50% of total blood volume is replaced within 3 hours

- The replacement of blood loss with red cells and a crystalloid volume expander will result in gradual dilution of plasma clotting proteins, leading to prolongation of the prothrombin time (PT) and the activated partial thromboplastin time (aPTT)

- A similar dilutional effect on the platelet concentration can be seen with massive transfusion

Massive transfusion protocol (MTPs)

- 
- Established to provide rapid blood replacement in a setting of severe haemorrhage
 - Early optimal blood transfusion is essential to sustain organ perfusion and oxygenation

Massive Transfusion-Clinical Settings


- Trauma
- Surgery (e.g. Liver, Cardiovascular)
- Less frequent
 - abdominal aortic aneurysm
 - liver transplant
 - obstetric catastrophes
 - GI bleeding

Challenges

- Types of components to be administered
- Selection of the appropriate amounts
- TIME



Emergency blood issue



**If units are issued without X match – written consent of physician to be taken,
– complete X match protocols followed after issue**

Haemostatic Resuscitation: FFP

- Meta-analysis from 2010-2012: Patients undergoing massive transfusion, high FFP to RBC ratios was associated with a significant reduction in the risk of death (odds ratio (OR) 0.38 (95%CI 0.24-0.60) and multiorgan failure (OR 0.40 (95%CI 0.26-0.60)).
- Murad MH, Stubbs JR, Gandhi MJ, Wang AT, Paul A, Erwin PJ, Montori VM, Roback JD: The effect of plasma transfusion on morbidity and mortality: a systematic review and meta-analysis. *Transfusion* 2010, 50:1370-1383

Haemostatic Resuscitation: FFP

- Meta-analysis from 2012 reports of reduced mortality in trauma patients treated with the highest FFP or PLT to RBC ratios.
- Johansson PI, Oliveri R, Ostrowski SR: Hemostatic resuscitation with plasma and platelets in trauma. *A meta-analysis. J Emerg Trauma Shock* 2012, 5:120-125.

Haemostatic Resuscitation: Plts

- Platelets are also pivotal for hemostasis: low Plts increases mortality.
- The highest survival was established in patients who received both a high PLT:RBC and a high FFP:RBC ratio.
- Holcomb JB, Wade CE, Michalek JE, Chisholm GB, Zarzabal LA, Schreiber MA, Gonzalez EA, Pomper GJ, Perkins JG, Spinella PC, Williams KL, Park MS: Increased plasma and platelet to red blood cell ratios improves outcome in 466 massively transfused civilian trauma patients. *Ann Surg* 2008, 248:447-458.

Recommendations

- “Damage control” approach
- Improved survival when the ratio of transfused Fresh Frozen Plasma (FFP, in units) to platelets (in units) to red blood cells (RBCs, in units) approaches 1:1:1

Holcomb JB, Jenkins D, Rhee P, et al. Damage control resuscitation: directly addressing the early coagulopathy of trauma. J Trauma 2007; 62:307.



Patients who have sustained severe traumatic injuries and/or who are likely to require massive transfusion should receive a 1:1:1 ratio of FFP to platelets to RBCs at the outset of their resuscitation and transfusion therapy

- Borgman MA, Spinella PC, Perkins JG, et al. The ratio of blood products transfused affects mortality in patients receiving massive transfusions at a combat support hospital. *J Trauma* 2007; 63:805.
- Holcomb JB, Wade CE, Michalek JE, et al. Increased plasma and platelet to red blood cell ratios improves outcome in 466 massively transfused civilian trauma patients. *Ann Surg* 2008; 248:447.
- Cotton BA, Au BK, Nunez TC, et al. Predefined massive transfusion protocols are associated with a reduction in organ failure and postinjury complications. *J Trauma* 2009; 66:41.
- Shaz BH, Dente CJ, Nicholas J, et al. Increased number of coagulation products in relationship to red blood cell products transfused improves mortality in trauma patients. *Transfusion* 2010; 50:493.
- Inaba K, Lustenberger T, Rhee P, et al. The impact of platelet transfusion in massively transfused trauma patients. *J Am Coll Surg* 2010; 211:573.
- de Biasi AR, Stansbury LG, Dutton RP, et al. Blood product use in trauma resuscitation: plasma deficit versus plasma ratio as predictors of mortality in trauma (CME). *Transfusion* 2011; 51:1925.

Fibrinogen concentrate

- European guidelines recommend fibrinogen concentrate when the level falls below 1.5g
- Cost of fibrinogen concentrate is much more than cryoprecipitate
- Availability






Cryoprecipitate

- Most common blood product used to replace fibrinogen
- Contains approximately 200–250 mg of fibrinogen per unit
- Standard dose of two 5-unit pools should be administered **early** in major **obstetric haemorrhage**.
- Subsequent **cryoprecipitate** transfusion should be guided by fibrinogen results, aiming to keep levels above 1.5 g/l.

Complications of Massive Transfusion

- Hypothermia
- Acid/base derangements
- Coagulopathy
- Citrate toxicity
- Electrolyte abnormalities
 - hypocalcemia
 - hypomagnesemia
 - hypokalemia
 - hyperkalemia
- Transfusion-associated acute lung injury



Acidosis and hypothermia

- Acidosis
 - Interferes with formation of coagulation factor complexes
- Hypothermia
 - Reduces enzymatic activity of coagulation factors
 - Prevents activation of platelets

Hypothermia

10 units of cold blood products and an hour of surgery can lead to a 3°C drop in core temperature and hypothermic coagulopathy

RBCs that are stored at 4C are transfused rapidly

Lowers the recipient's core temperature and further impairs haemostasis.

Reduces the metabolism of citrate and lactate
Increases the likelihood of hypocalcaemia, metabolic acidosis and cardiac arrhythmias.

Shifts the oxyhaemoglobin dissociation curve to the left, reducing tissue oxygen delivery

Prevention of hypothermia

- A high capacity commercial blood warmer should be used to warm blood components



Coagulopathy

- Dilutional coagulopathy
- Disseminated intravascular coagulation.
- Consumption of platelets and coagulation factors

- 500 mL blood loss replaced → 10% drop in clotting factor activity
- 8 – 10 units of PRBCs → coagulation activity at 25%

ALTERATIONS IN HEMOSTASIS

- Acute DIC
 - microvascular oozing
 - prolongation of the PT and aPTT in excess of that expected by dilution
 - significant thrombocytopenia
 - low fibrinogen levels
 - increased levels of D-dimer

Hypocalcaemia

- Citrate binds calcium
- Results in hypotension, small pulse pressure, flat ST-segments and prolonged QT intervals on the ECG.
- Slow i.v. injection of calcium gluconate 10%



Hyperkalaemia

- The potassium concentration of blood increases during storage, by as much as 5–10 mmol u1 .
- Hyperkalaemia rarely occurs during massive transfusions unless the patient is also hypothermic and acidotic

Monitoring recommendations

- PT, aPTT
- Platelet count
- Fibrinogen
- Electrolytes
- Viscoelastic test
 - after the administration of every five to seven units of red cells.

Goals


Massive Blood Transfusion Management

- Haemostatic Resuscitation
- Fluid management
- Metabolic acid base correction
- Normal temperature
- Calcium management
- Management of Coagulopathy

Viscoelastic whole-blood assays








- TEG[®] and ROTEM[®]
- provide information on the coagulation process through the graphic display of clot initiation, propagation and lysis.
- used to guide transfusion of blood components

Measure all phases of hemostasis in whole blood.



The TEG® hemostasis system continuously measures all phases of hemostasis as a net product of whole blood components

- Costeffective -since it reduces inappropriate transfusions, thus improving transfusion management and patients' clinical outcome

Laboratory Value	Interpretation	Blood Product Transfusion	QUALITATIVE INTERPRETATION — PATTERN RECOGNITION
R less than 4 min	Enzymatic Hypercoagulability	No treatment if bleeding	 Normal R, K, MA, Angle Normal
R between 11-14 min	Low clotting factors	Plasma and RBC's	 Anticoagulant/hemophilia Factor Deficiency R, K = Prolonged MA = Decreased
R greater than 14 min	Very low clotting factors	Plasma and RBC's	 Platelet Blockers Thrombocytopenia/Thrombotic R = Normal, K = Prolonged MA = Decreased
a-angle < 45 degrees	Low fibrinogen level	Cryoprecipitate/ Fibrinogen /Platelets	 Fibrinolysis R = Normal MA = Continuous Decrease
MA between 46-54 mm	Low platelet function	Platelets/ Cryoprecipitate/ Fibrinogen	 Hypercoagulation R, K = Decreased MA = Increased
MA between 41-45 mm	Very low platelet function	Platelets/ Cryoprecipitate/ Fibrinogen	 D.I.C. Stage 1 - Hypercoagulable + secondary fibrinolysis
MA at 40 mm or less	Extremely very low platelet function	Platelets/ Cryoprecipitate/ Fibrinogen	 Stage 2 - Hypocoagulable state
MA greater than 73 mm	Platelet Hypercoagulability	No treatment if bleeding	
LY30 greater than 3%, CI less than 1.0	Primary fibrinolysis	Tranexamic acid 1g IV over 10 minutes followed by 1g in 250cc NS infused over 8 hours	

• Refer to TEG analysis tree for values outside these ranges

Thank You