



Cardiac Arrhythmias

KAIS AL BALBISSI, MD, FACC, FSCAI

ASSOCIATE PROFESSOR OF INTERNAL MEDICINE





Cardiac Arrhythmias



Cardiac Arrhythmias

- ECG Basics
- Approach to Rhythm
- Brady-Arrhythmias
 - Sinus Bradycardia
- Arrhythmias
 - Premature Atrial Contraction
 - Premature Ventricular Contraction
- Tachy-Arrhythmias
 - Sinus Tachycardia
 - Atrial Fibrillation
 - Atrial Flutter
 - Multifocal Atrial Tachycardia
 - Supraventricular Tachycardias
 - Ventricular Tachyarrhythmias

ECG Basics





ECG Basics

- **Approach to Reading ECG:**

- Verify
- Rate
- Rhythm
- Axis
- Amplitude
- Intervals
- Ischemia

ECG Basics



The whole ECG Strip is 10 seconds

ECG Basics - Rate

- Heart Rate Calculation Methods:
 - Counts QRS complexes
 - 6 second interval X 10
 - All strip (10 seconds) X 6
 - Distance between QRS-QRS complex
 - Rate = $300 / \text{Large Boxes}$
 - Rate = $1500 / \text{Small Boxes}$




# Large Boxes	HR (BPM)
1	300
2	150
3	100
4	75
5	60
6	50

ECG Basics - Rhythm

Approach to Rhythm Questions:	Clinical Significance
1. Is it Tachycardia / Normal Rate / Bradycardia?	Rate
2. QRS is it Narrow or Wide?	<u>Narrow</u> : Rhythm from AVN and above & conduction through normal system <u>Wide</u> : Rhythm below AVN OR Abnormal conduction
3a. Narrow QRS – Is it Regular or Irregular	
3b. Wide QRS – What is Morphology?	Pathophysiology of Wide QRS: Vent. Origin or Aberrant conduction?
4. Look for P-wave (Best place in Lead II and V1)	What is the atria doing?
5. Relationship between the P wave and QRS ?	What is the underlying circuit?


ECG Basics - Axis



ECG Basics - Axis


- Heart Axis Calculation Methods:
 - Lead I, II & III

Axis	Lead I	Lead II	Lead III
Normal	Positive	Positive	Positive
LAD	Positive	Negative	Negative
RAD	Negative	Positive	Positive
Extreme Axis	Negative	Negative	Negative



ECG Basics - Axis

- Heart Axis Calculation Methods:
 - Lead I & Lead AVF




Copyright ©2006 by The McGraw-Hill Companies, Inc.
All rights reserved.

ECG Basics - Axis

Extreme
Axis





- Heart Axis Calculation Methods:
 - Lead I, II & III

Axis	Lead I	Lead II	Lead III
Normal	Positive	Positive	Positive
LAD	Positive	Negative	Negative
RAD	Negative	Positive	Positive
Extreme Axis	Negative	Negative	Negative



ECG Basics - Axis

12-lead ECG Electrode Placement



ECG Basics – Amplitude / Hypertrophy


- Low Amplitude

- Limb leads < 0.5 mV
- Precordial leads < 1.0 mV

Component	Amplitude (mV)
P wave	0.2
QRS	1.0
T-wave	0.2 - 0.3


ECG Basics - Amplitude / Hypertrophy

RAE



© 1997 Frank G. Yanowitz, M.D.

LAE



© 1997 Frank G. Yanowitz, M.D.

P wave amplitude > 2.5 mm in II and/or > 1.5 mm in V₁

P wave duration ≥ 0.12s in frontal plane (usually lead II)
Notched P wave in limb leads with the inter-peak duration ≥ 0.04s
Terminal P negativity in lead V₁ (i.e., "P-terminal force") duration ≥ 0.04s
& depth ≥ 1 mm.

ECG Basics – Amplitude / Hypertrophy

LVH

ESTES Criteria

- Voltage Criteria (any of): R or S in limb leads ≥ 20 mm
- S in V1 or V2 ≥ 30 mm
- R in V5 or V6 ≥ 30 mm

- ST-T Abnormalities: Without digitalis
- With digitalis

Left Atrial Enlargement in V1

Left axis deviation

QRS duration 0.09 sec

Delayed intrinsicoid deflection in V5 or V6 (>0.05 sec)

Points

3 points

3 points

1 point

3 points

2 points

1 point

1 point

("diagnostic", ≥ 5 points; "probable", 4 points)

CORNELL Voltage Criteria

- S in V3 + R in aVL > 24 mm (men)
- S in V3 + R in aVL > 20 mm (women)

(sensitivity = 22%, specificity = 95%)

Limb-lead voltage criteria:

- R in aVL ≥ 11 mm
- R in aVL ≥ 13 mm + S in III ≥ 15 mm (if LAD)
- R in I + S in III > 25 mm

Chest-lead voltage criteria:


- S in V1 + R in V5 or V6 ≥ 35 mm

ECG Basics - Amplitude / Hypertrophy

RVH

- Any one or more of the following (if QRS duration < 0.12 sec):
 - Right axis deviation (> 90 degrees) in presence of disease capable of causing RVH
 - R in aVR ≥ 5 mm, or
 - R in aVR > Q in aVR
- Any one of the following in lead V1:
 - R/S ratio > 1 and negative T wave
 - qR pattern
 - R > 6 mm, or S < 2mm, or rSR' with R' > 10 mm
- Other chest lead criteria:
 - R in V1 + S in V5 (or V6) 10 mm
 - R/S ratio in V5 or V6 < 1
 - R in V5 or V6 < 5 mm
 - S in V5 or V6 > 7 mm


ECG Basics - Intervals




$$QTc = QT \text{ Interval} / \sqrt{RR \text{ Interval}}$$

Upper Limit of Normal QTc	ms
Male	> 460 - 470
Female	> 470 - 480

ECG Basics - Ischemia



ECG Basics - Ischemia



Approach to ECG

▪ Rate


- Regular: Rate = 300 / Large Boxes
- Irregular: Rate = # R in ECG X 6

▪ Rhythm

Approach to Rhythm Questions:

1. Is it Tachycardia / Normal Rate / Bradycardia?
2. QRS is it Narrow or Wide?
- 3a. Narrow QRS – Is it Regular or Irregular
- 3b. Wide QRS – What is Morphology?
4. Look for P-wave (Best place in Lead II and V1)
5. Relationship between the P wave and QRS ?

▪ Axis



▪ Amplitude

Low:

Limb < 0.5 mm
Chest < 1.0 mm

IAE:

P Width > 120ms

RAE:

P Ht. > 2.5 mm

LVH : Cornell's Criteria

- S in V3 + R in aVL > 24 mm (men)
- S in V3 + R in aVL > 20 mm (women)

LVH: Lead AVL > 11 mm

RVH : Lead V1

- R/S ratio > 1 and negative T wave
- R > 6 mm / S < 2mm
- rSR' with R' > 10 mm


▪ Intervals

Intervals	# Small sq.
PR	120-200 ms
QRS	< 110-120 ms
QT	< 480-500 ms

$< \frac{1}{2}$ RR Interval
< 12

▪ Ischemia


STEMI



Non - ST ACS



Approach to Rhythm



Approach to Rhythm - Normal Rate

Question	Answer								
1. Rate	Normal Rate								
2. QRS	Narrow								
3. QRS	Regular					Irregular			
								Morphology	
								Ventricular	Aberrancy
4. P wave	Present			Absent	Present	Absent	Absent	Present	
5. P-QRS Relation	Single	Multiple	Retrograde						
DDx	NSR	A.Flutter	Junctional Rhythm	Junctional Rhythm	SR with PAC	A.Fib	AIVR	Conduction Abnormality	
	Atrial Rhythm	2:1 AV Block			Wandering Pacemaker				
	1 st degree AV Block				A.FI with Variable Conduction				

Approach to Rhythm - Bradycardia

Question	Answer										
1. Rate	Bradycardia										
2. QRS	Narrow										
3. QRS	Regular										
4. P wave	Present						Absent	Present	Absent	Absent	Present
5. P-QRS Relation	Single	Multiple	Group Beating	Retrograde	AV Dissociation						
DDx	SB	A.Flutter with SVR	2 nd degree AV Block	Junctional Escape Rhythm	Complete AV Block (3 rd degree)	Junctional Escape Rhythm	SB with PAC	A.Fib with SVR	Ventricular Escape Rhythm	Conduction Abnormality	
	1st degree AV Block	2:1 AV Block					A.Fi with Slow & Variable Conduction				

Approach to Rhythm - Tachycardia

- DDx of SVT

- Short RP Tachycardias (RP<PR):

- Typical AV Nodal Re-entry Tachycardia (AVNRT)
 - Junctional Tachycardia
 - Orthodromic Atrioventricular Tachycardia (OD – AVRT)
 - Atrial Tachycardia


Short R-P



- Long RP Tachycardias (RP>PR):

- Sinus Tachycardia (ST)
 - Atrial Tachycardia (AT)
 - Atypical Orthodromic Atrioventricular Tachycardia (OD – AVRT)
 - Atypical AV Nodal Re-entry Tachycardia (AVNRT)
 - Junctional Tachycardia

Long R-P



- Mimickers:


- Atrial Flutter with rapid conduction
 - A. Fibrillation with very rapid conduction

Approach to Rhythm - Tachycardia


Tachyarrhythmia Framework

	<u>REGULAR RHYTHM</u>	<u>IRREGULAR RHYTHM</u>
NARROW QRS	<ul style="list-style-type: none">• Sinus tachycardia• AVNRT• Orthodromic AVRT• Atrial Tachycardia• Atrial flutter• Junctional tachycardia	<ul style="list-style-type: none">• Atrial fibrillation• Atrial flutter w/ variable AV block• Multifocal atrial tachycardia
WIDE QRS	<ul style="list-style-type: none">• Ventricular tachycardia• SVT with bundle branch block• Antidromic AVRT• Pre-excited SVT	<ul style="list-style-type: none">• Polymorphic ventricular tachycardia• Atrial fibrillation with bundle branch block• Atrial flutter with variable AV block & bundle branch block

Arrhythmias



Mechanisms of Arrhythmias



Arrhythmias


- Premature Atrial Contraction / Complex
- Premature Ventricular Contraction / Complex

Premature Atrial Contractions / Complexes (PAC)

- Very common. May cause Palpitations

- Causes:



- Adrenergic excess
- Pharmacological
- Electrolyte imbalances
- Ischemia
- Hypoxia
- Infection.



Clinical Status	Management
Asymptomatic	Observation
Symptomatic (Palpitations, Fatigue, Exercise Intolerance, Angina, Dizziness, Syncope)	Rx Cause B-Blockers

Premature Ventricular Contractions / Complexes (PVC)

- Common. May cause Palpitations.
- Causes:
 - Hypoxia
 - Electrolyte abnormalities
 - Pharmacological
 - Structural heart disease




Copyright © 2015 Lippincott

Premature Ventricular Contractions / Complexes (PVC)

Premature Ventricular Contraction (PVC) - Subtypes





BIGEMINY



TRIGEMINY



QUADRIGEMINY




NSVT

Premature Ventricular Contractions / Complexes (PVC)

Clinical Status	Management
Asymptomatic - Infrequent	Observation
Asymptomatic - Frequent / Repetitive	R/o Heart Disease B-Blockers EPS +/- ICD/Ablation
Symptomatic (Palpitations, Fatigue, Exercise Intolerance, Angina, Dizziness, Syncope)	Rx Cause B-Blockers

Tachyarrhythmias




Tachyarrhythmias

- Sinus Tachycardia
 - Inappropriate Sinus Tachycardia
- Atrial Fibrillation
- Atrial Flutter
- Multifocal Atrial Tachycardia
- Supraventricular Tachycardia
- Ventricular Tachycardia
- Ventricular Fibrillation

Sinus Tachycardia

- Rate >100 bpm

Sinus Tachycardia



shutterstock.com • 1835823070

Sinus Tachycardia

- Causes:



Physiological	Cardiac Conditions	Medical Condition	Pharmacological
<ul style="list-style-type: none">• Exercise• Emotion• Anxiety• Pain• Fever• Pregnancy• <i>Volume Depletion</i>	<ul style="list-style-type: none">• MI• Cardiomyopathy / HF• Acute Valve Disease• Pericarditis• Postural	<ul style="list-style-type: none">• Shock• Hypoxia• Respiratory Distress• P.E.• Anemia• Infection• <i>Dehydration</i>• Hyperthyroidism• Pheochromocytoma• Cushing's• Hypoglycemia• Panic Attack	<ul style="list-style-type: none">• Caffeine• Alcohol• Tobacco• Catecholamines• B agonist• BB Withdrawal• Vasodilators• Atropine• Theophylline• Decongestants• Cocaine• Amphetamines• Thyroid Hormones

Sinus Tachycardia

- Management:
 - Treat the underlying cause
 - Treat the underlying cause
 - Treat the underlying cause
- Inappropriate Sinus Tachycardia
 - B-Blockers
 - Ivabradine
 - RFA

Atrial Fibrillation

- Very common



Atrial Fibrillation

- Causes:
 - Heart disease: CAD, MI, HTN, mitral valve disease
 - History of cardiac surgery
 - Pericarditis
 - Pulmonary disease (PE, COPD, Hypoxia)
 - Thyroid disease
 - Pheochromocytoma
 - Systemic illness (e.g. Infection,)
 - Stress (postoperative, pain, anxiety)
 - Hyperadrenergic states
 - Cocaine or methamphetamine use
 - Extremes of activity (sedentary lifestyle, excess exercise such as marathon running)
 - Excessive alcohol intake (“holiday heart syndrome”)

Atrial Fibrillation - Management

Rate Control

B-Blockers

CCB

Digoxin

Anticoagulation

Warfarin

DOACS

CHA2DS2-VASc Score

CHF or LVEF ≤ 40%	1
HTN	1
Age ≥ 75	2
DM	1
CVA/TIA/TE	2
Vascular	1
Age 65-74	1
Female	1

HAS-BLED score

Condition	Points
H - Hypertension	1
A - Abnormal renal or liver function (1 point each)	1 or 2
S - Stroke	1
B - Bleeding	1
L - Labile INRs	1
E - Elderly (> 65 years)	1
D - Drugs or alcohol (1 point each)	1 or 2

Atrial Fibrillation - Management

Rate Control

B-Blockers

CCB

Digoxin

Anticoagulation

Warfarin

DOACS

CHA2DS2-VASc Score

CHF or LVEF ≤ 40%	1
H - Hypertension	1
CVA Risk	
A 0	0
D 1	1.3
C 2	2.2
V 3	3.2
V 4	4
A 5	6.7
F 6	9.8
7	9.6
8	6.7
9	15.2

HAS-BLED score

Condition	Points
H - Hypertension	1
A - Abnormal renal or liver function (1 point each)	1 or 2
S - Stroke	
B - Bleeding	
L - Labile INR	
E - Elderly	0 1.13
D - Drug	1 1.02
	2 1.88
	3 3.74
	4 8.70
	5 12.5

Atrial Fibrillation - Pearls

The image shows two screenshots of a mobile application interface for managing atrial fibrillation (AF). The top navigation bar includes 'Sprint' signal, '10:13 AM', and '96%'. The main menu has 'Calculate Risk' and 'Review Therapy' buttons.

Left Screenshot (Calculate Risk):

- Stroke Risk:** CHA₂DS₂-VASc score: 3 (High risk).
- Renal Function:** eGFR: 60 mL/min (Normal).

Right Screenshot (Review Therapy):

- Stroke Risk:** CHA₂DS₂-VASc score: 3 (High risk).
- Bleeding Risk:** 1.2 mg/dL INR (Low risk) and 61.1 mL/min eGFR (Normal).

Calculate Risk and **Reset All** buttons are present at the bottom of the left screen.

Patient Information: Required to derive therapy options.

Age: [Input field] Yrs

Sex: [Dropdown menu] Please select

CHA₂DS₂-VASc: Select all that apply.

CHF/LV dysfunction

② Select Therapy Option: Dabigatran

③ Evaluate Therapy:

Standard Dose: 150 mg twice daily (clinical trials)

Risk/Benefit Information*

Patient's ANNUAL risk of stroke + thromboembolism with Dabigatran	0.9%
Relative risk reduction	79%
Absolute risk reduction	3.4%
Chance of benefit per year	1 in 30

Atrial Fibrillation - Management

Rate Control

B-Blockers

CCB

Digoxin

Anticoagulation

Warfarin

DOACS

CHA2DS2-VASc Score

CHF or LVEF ≤ 40%	1
HTN	1
Age ≥ 75	2
DM	1
CVA/TIA/TE	2
Vascular	1
Age 65-74	1
Female	1

If

Unstable

Symptomatic

Rate not controlled

Young / New Onset



Rhythm Control

Electrical

Pharmacological


Ablation

Atrial Fibrillation - Pearls




Atrial Flutter

- Common





Atrial Flutter - Management




Atrial Flutter - Pearls

Atrial Flutter



Multifocal Atrial Tachycardia (MAT)


- Common in patients with severe pulmonary disease (e.g., COPD)



Multifocal Atrial Tachycardia (MAT)

- Treatment:
 - Improve oxygenation and ventilation
 - Medications: CCBs, β -blockers, digoxin, amiodarone
 - Electrical cardioversion is ineffective and should not be used

Supraventricular Tachycardia (SVT)




Supraventricular Tachycardia (SVT)

- DDx of SVT


- Short RP Tachycardias (RP<PR):

- Typical AV Nodal Re-entry Tachycardia (AVNRT)
 - Junctional Tachycardia
 - Orthodromic Atrioventricular Tachycardia (OD – AVRT)
 - Atrial Tachycardia



- Long RP Tachycardias (RP>PR):



- Sinus Tachycardia (ST)
 - Atrial Tachycardia (AT)
 - Atypical Orthodromic Atrioventricular Tachycardia (OD – AVRT)
 - Atypical AV Nodal Re-entry Tachycardia (AVNRT)
 - Junctional Tachycardia





- Mimickers:

- Atrial Flutter with rapid conduction
 - A. Fibrillation with very rapid conduction

SVT: AV Nodal Re-entrant Tachycardia (AVNRT)





SVT: AV Nodal Re-entrant Tachycardia (AVNRT)





Management:

- Acute management:**
 - Vagal Maneuvers
 - Pharmacologic therapy
 - IV adenosine
 - CCB / BB
 - DCCV
- Preventative management:**
 - Pharmacologic therapy: CCBs or β -blockers
 - Ablation

SVT: Orthodromic AV Re-entrant Tachycardia (AVRT)





SVT: Orthodromic AV Re-entrant Tachycardia (AVRT)



Management:



- Acute management:
 - Vagal Maneuvers
 - Pharmacologic therapy
 - IV adenosine
 - CCB / BB
 - DCCV
- Preventative management:
 - Pharmacologic therapy: CCBs or β -blockers
 - Ablation

SVT: Antidromic AV Re-entrant Tachycardia (AVRT)



SVT: Wolf Parkinson White Syndrome (WPW)

Pre-excitation





SVT: Wolf Parkinson White Syndrome (WPW)

Pre-excitation

+

SVT

= WPW





SVT: Wolf Parkinson White Syndrome (WPW)

Pre-excitation

+

SVT

= WPW



Management:

- Acute management:
 - Pharmacologic therapy
 - Procainamide
 - DCCV
- Preventative management:
 - Ablation
- Do not use CCBs or β -blockers or Digoxin


Ventricular Tachycardia (VT)

- **Causes:**
 - Ischemia
 - CAD with prior MI is the most common cause
 - Cardiomyopathies
 - Ventricular scar tissue
 - Congenital defects
 - Long QT syndrome
 - Electrolyte Abnormalities
 - Drug toxicity (antiemetics, antipsychotics, SSRIs, TCAs, macrolide and fluoroquinolone antibiotics)


Ventricular Tachycardia (VT)

- Non-Sustained Ventricular Tachycardia (NSVT): < 30 seconds

Monomorphic




Polymorphic




Ventricular Tachycardia (VT)

- Sustained Ventricular Tachycardia (NSVT): > 30 seconds

Monomorphic



Polymorphic



Torsades de Points (TdP)


Stable vs. Unstable?

Ventricular Tachycardia (VT) - Stable

Management:

- Acute management:
 - Treat the underlying cause:
 - Ischemia
 - Correct Electrolyte Abnormalities
 - Remove Drug +/- Antidote
 - Pharmacological Therapy:
 - IV Amiodarone
 - DCCV
- Preventative management:
 - Consider ICD
 - Consider EPS

Ventricular Vibration (VF)



Code Blue
Defibrillation

& Unstable Sustained VT

Bradyarrhythmias




Bradycardia

- Sinus Bradycardia
- Sick Sinus Syndrome (SSS)
- Atrio-Ventricular (AV) Block
 - 1st Degree
 - 2nd Degree
 - Mobitz Type I (Wenckebach)
 - Mobitz Type II
 - 2:1 Block
 - 3rd Degree (Complete)
- Pacemakers & Cardiac Devices

Sinus Bradycardia

- Rate < 60 bpm
- Causes:
 - Ischemia
 - Increased Vagal tone
 - Structural Heart Disease (Infiltrative, IE, ACHD)
 - Medications
 - Athletes



Clinical Status

Asymptomatic

Symptomatic
(Fatigue, Exercise Intolerance, Angina, Dizziness, Syncope)


Management

Observation

Rx Cause
Atropine / B Agonist (Acute Rx)
Pacemaker

Sick Sinus Syndrome (SSS)

- SA Dysfunction



- Management:
 - Pacemaker placement

Occurs with advanced age
Marked Persistent Sinus Bradycardia
SA Pauses and Blocks
Frequently associated with Tachy-Brady Syndrome
Usually co-exists with AV nodal disease

Atrio-Ventricular (AV) Block

- 1st Degree AV Block
 - Prolonged PR Interval > 0.2 second
 - No dropped beats (*No P without QRS*)

Causes:

- Ischemia
- Increased Vagal tone
- Structural Heart Disease (Infiltrative, IE, ACHD)
- Medications




- Management:

Clinical Status	Management
Asymptomatic	Observation
Symptomatic (Fatigue, Exercise Intolerance, Angina, Dizziness, Syncope)	Rx Cause Atropine / B Agonist (Acute Rx) Pacemaker

Atrio-Ventricular (AV) Block

- 2nd Degree AV Block – Mobitz Type I (Wenckebach)
 - Progressive PR Prolongation followed by a dropped QRS

- Causes:
- Ischemia
 - Increased Vagal tone
 - Structural Heart Disease
 - Medications




- Management:

Clinical Status	Management
Asymptomatic	Observation
Symptomatic (Fatigue, Exercise Intolerance, Angina, Dizziness, Syncope)	Rx Cause Atropine / B Agonist (Acute Rx) Pacemaker

Atrio-Ventricular (AV) Block

- **2nd Degree AV Block – Mobitz Type II**
 - Progressive PR Prolongation followed by a dropped QRS



- Management:
 - Pacemaker placement indicated

Atrio-Ventricular (AV) Block

- **2nd Degree AV Block (2:1 Block)**
 - Alternating conducted QRS followed by a dropped QRS



- **Management:**
 - Needs further evaluation



Atrio-Ventricular (AV) Block

- **3rd Degree AV Block**
 - P > QRS
 - AV Dissociation
- **Management:**
 - Medical Emergency
 - Emergent Pacer placement




Atrio-Ventricular (AV) Block

- 3rd Degree AV Block



Atrio-Ventricular (AV) Block

- 3rd Degree AV Block




Pacemakers & Cardiac Device

Temporary

Transcutaneous





Transvenous (TVP)




Pacemakers & Cardiac Devices

Permanent Pacemaker (PPM)





Letter 1	Letter 2	Letter 3	Letter 4	Letter 5
Chamber Paced	Chamber Sensed	Sensing Response	Programmability	Anti-tachycardia Functions
A = Atrium	A = Atrium	T = Triggered	P = Simple	P = Pacing
V = Ventricle	V = Ventricle	I = Inhibited	M = Multiprogrammable	S = Shock
D = Dual	D = Dual	D = Dual (Inhibits Both the Atrium & Ventricle)	R = Rate Adaptive	D = Dual (Shock Pace)
O = None	O = None	O = None	O = None	





Pacemakers & Cardiac Devices

Implantable Cardiac Defibrillator (ICD)



Pacemakers & Cardiac Devices

Cardiac Resynchronization Therapy (CRT)



AKA
Bi Ventricular Pacer
(BiV PPM)

Types	
CRT - P	CRT - D
CRT only	CRT with Defibrillator

Pacemakers & Cardiac Devices






Figure 1: 174105-2 Magnet

Magnet Mode

	Magnet Mode ON
Pacemaker	Asynchronous Pacing
ICD	Defibrillator OFF

Cardiac Arrhythmias



Thank You